

## TABLE OF CONTENTS

### SPOTLIGHT

|  |    |
|--|----|
| Articles of Significant Interest Selected from This Issue by the Editors | 89 |
|--|----|

### COMMENTARY

|  |  |       |
|--|--|-------|
| The State of Research for AIDS-Associated Opportunistic Infections and the Importance of Sustaining Smaller Research Communities | Anthony P. Sinai, Edna S. Kaneshiro, Honorine Ward, Louis M. Weiss, and Melanie T. Cushion | 90–97 |
|--|--|-------|

### MINIREVIEWS

|  |  |         |
|--|--|---------|
| Posttranslational Modifications of Proteins in the Pathobiology of Medically Relevant Fungi                        | Michelle D. Leach and Alistair J. P. Brown   | 98–108  |
| Adaptation of <i>Cryptococcus neoformans</i> to Mammalian Hosts: Integrated Regulation of Metabolism and Virulence | Jim Kronstad, Sanjay Saikia, Erik David Nielson, Matthias Kretschmer, Wonhee Jung, Guanggan Hu, Jennifer M. H. Geddes, Emma J. Griffiths, Jaehyuk Choi, Brigitte Cadieux, Mélissa Caza, and Rodgoun Attarian | 109–118 |

### ARTICLES

|   |  |         |
|---|--|---------|
| Protein Kinase A Contributes to the Negative Control of Snf1 Protein Kinase in <i>Saccharomyces cerevisiae</i>  | LaKisha Barrett, Marianna Orlova, Marcin Maziarz, and Sergei Kuchin  | 119–128 |
| Transcription Factor Efg1 Shows a Haploinsufficiency Phenotype in Modulating the Cell Wall Architecture and Immunogenicity of <i>Candida albicans</i> | Martin Zavrel, Olivia Majer, Karl Kuchler, and Steffen Rupp  | 129–140 |
| Heterogeneous Expression of the Virulence-Related Adhesin Epa1 between Individual Cells and Strains of the Pathogen <i>Candida glabrata</i>           | Samantha C. Halliwell, Matthew C. A. Smith, Philippa Muston, Sara L. Holland, and Simon V. Avery   | 141–150 |
| MADS Box Transcription Factor Mbx2/Pvg4 Regulates Invasive Growth and Flocculation by Inducing <i>gsf2</i> <sup>+</sup> Expression in Fission Yeast   | Tomohiko Matsuzawa, Ken-ichi Yoritsune, and Kaoru Takegawa   | 151–158 |
| Snf1-Like Protein Kinase Ssp2 Regulates Glucose Derepression in <i>Schizosaccharomyces pombe</i>  | Tomohiko Matsuzawa, Yasuko Fujita, Hideki Tohda, and Kaoru Takegawa  | 159–167 |
| Rhb1 Regulates the Expression of Secreted Aspartic Protease 2 through the TOR Signaling Pathway in <i>Candida albicans</i>                            | Yu-Ting Chen, Chia-Ying Lin, Pei-Wen Tsai, Cheng-Yao Yang, Wen-Ping Hsieh, and Chung-Yu Lan  | 168–182 |
| Enigmatic Presence of Mitochondrial Complex I in <i>Trypanosoma brucei</i> Bloodstream Forms  | Sachin Surve, Meredith Heestand, Brian Panicucci, Achim Schnauffer, and Marilyn Parsons  | 183–193 |
| The Polo-Like Kinase PLKA in <i>Aspergillus nidulans</i> Is Not Essential but Plays Important Roles during Vegetative Growth and Development          | Klarita Mogilevsky, Amandeep Glory, and Catherine Bachewich  | 194–205 |
| SPM1 Stabilizes Subpellicular Microtubules in <i>Toxoplasma gondii</i>  | Johnson Q. Tran, Catherine Li, Alice Chyan, Lawton Chung, and Naomi S. Morrisette  | 206–216 |
| A Machine Learning Approach To Identify Hydrogenosomal Proteins in <i>Trichomonas vaginalis</i>   | David Burstein, Sven B. Gould, Verena Zimorski, Thorsten Kloesges, Fuat Kiosse, Peter Major, William F. Martin, Tal Pupko, and Tal Dagan | 217–228 |

Continued on following page

**SWP5, a Spore Wall Protein, Interacts with Polar Tube Proteins in the Parasitic Microsporidian *Nosema bombycis***

Zhi Li, Guoqing Pan, Tian Li, Wei Huang, Jie Chen, Lina Geng, Donglin Yang, Linling Wang, and Zeyang Zhou 229–237

**Resolving Phenylalanine Metabolism Sheds Light on Natural Synthesis of Penicillin G in *Penicillium chrysogenum***

Tânia Veiga, Daniel Solis-Escalante, Gabriele Romagnoli, Angela ten Pierick, Mark Hanemaaijer, Amit Deshmukh, Aljoscha Wahl, Jack T. Pronk, and Jean-Marc Daran 238–249

**GENOME ANNOUNCEMENT**

**Genome Sequence of the Fungus *Glarea lozoyensis*: the First Genome Sequence of a Species from the *Helotiaceae* Family**

Loubna Youssar, Björn Andreas Grüning, Anika Erxleben, Stefan Günther, and Wolfgang Hüttel 250

---

Cover photograph (Copyright © 2012, American Society for Microbiology. All Rights Reserved.): Localization of SPM1 to *Toxoplasma gondii* tachyzoite subpellicular microtubules. Stable cortical microtubules and the associated conoid were isolated by extraction in deoxycholate, and the resulting sample was stained with a *Toxoplasma*-specific tubulin antibody (red) and mouse antiserum that recognizes SPM1 (green). SPM1 localizes to the full length of the subpellicular microtubules but not the conoid which is located at the intersection of the subpellicular microtubules. (See related article on page 206.)