

# Regulation of Ammonia Homeostasis by the Ammonium Transporter AmtA in *Dictyostelium discoideum*<sup>∇</sup>

Ryuji Yoshino,<sup>1,5\*</sup> Takahiro Morio,<sup>1</sup> Yoko Yamada,<sup>2,†</sup> Hidekazu Kuwayama,<sup>1,3</sup>  
 Masazumi Sameshima,<sup>2,4</sup> Yoshimasa Tanaka,<sup>1</sup>  
 Hiromi Sesaki,<sup>5</sup> and Miho Iijima<sup>5\*</sup>

Graduate School of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Ibaraki, Japan 305-8572<sup>1</sup>;  
 Electron Microscopy Center, The Tokyo Metropolitan Institute of Medical Science, Bunkyo-ku, Tokyo,  
 Japan 113-8613<sup>2</sup>; Graduate School of Frontier Biosciences, Osaka University, Suita, Osaka,  
 Japan 565-0871<sup>3</sup>; Department of Biofunctional Science Faculty of Agriculture and Life Science,  
 Hirosaki University, Bunkyo-cho, Hirosaki, Aomori, Japan 036-8561<sup>4</sup>; and  
 Department of Cell Biology, Johns Hopkins University, School of  
 Medicine, Baltimore, Maryland 21205<sup>5</sup>

Received 11 June 2007/Accepted 3 October 2007

**Ammonia has been shown to function as a morphogen at multiple steps during the development of the cellular slime mold *Dictyostelium discoideum*; however, it is largely unknown how intracellular ammonia levels are controlled. In the *Dictyostelium* genome, there are five genes that encode putative ammonium transporters: *amtA*, *amtB*, *amtC*, *rhgA*, and *rhgB*. Here, we show that AmtA regulates ammonia homeostasis during growth and development. We found that cells lacking *amtA* had increased levels of ammonia/ammonium, whereas their extracellular ammonia/ammonium levels were highly decreased. These results suggest that AmtA mediates the excretion of ammonium. In support of a role for AmtA in ammonia homeostasis, AmtA mRNA is expressed throughout the life cycle, and its expression level increases during development. Importantly, AmtA-mediated ammonia homeostasis is critical for many developmental processes. *amtA*<sup>−</sup> cells are more sensitive to NH<sub>4</sub>Cl than wild-type cells in inhibition of chemotaxis toward cyclic AMP and of formation of multicellular aggregates. Furthermore, even in the absence of exogenously added ammonia, we found that *amtA*<sup>−</sup> cells produced many small fruiting bodies and that the viability and germination of *amtA*<sup>−</sup> spores were dramatically compromised. Taken together, our data clearly demonstrate that AmtA regulates ammonia homeostasis and plays important roles in multiple developmental processes in *Dictyostelium*.**

The cellular slime mold *Dictyostelium discoideum* has a unique life cycle consisting of a unicellular growth phase and a multicellular developmental phase. When food sources such as bacteria are available, *Dictyostelium* amoeboid cells proliferate by cytokinesis. Starvation triggers cells to undergo developmental processes, during which up to 10<sup>5</sup> cells display chemotaxis toward cyclic AMP (cAMP) and form multicellular aggregates. On top of the aggregates, a small projection is formed, and this process is called tip formation. Cells located at the anterior part of aggregates differentiate into prestalk cells, precursors of stalk cells, while the rest of the aggregates become prespore cells, precursors of spores. The tipped aggregates form elongated multicellular structures called slugs. Slugs migrate and eventually culminate to form a fruiting body consisting of a mass of spores supported by a stalk (25, 44, 47).

A number of diffusible molecules regulate the development of *Dictyostelium*, including cAMP, differentiation-inducing factor, adenosine, and ammonia (5, 47, 74, 78). Ammonia has

been shown to affect many developmental events in *Dictyostelium*. For example, in the presence of ammonia, the production and secretion of cAMP are inhibited, resulting in the impairment of chemotaxis toward cAMP and subsequent tip formation during early development (23, 61, 77). At later stages of development, ammonia acts against differentiation-inducing factor, suppresses differentiation into prestalk cells, and promotes differentiation into prespore cells (6, 26, 64, 72). In addition, ammonia plays an important role in the choice between the formation of a migrating slug and culmination. High concentrations of ammonia keep slugs migrating and block the initiation of culmination (60). The exhaustion of the ammonia supply triggers culmination at least in part by activating protein kinase A (PKA) through the DhkC signaling pathway (30, 61, 63). In fruiting bodies, extremely high concentrations of ammonium phosphate in sori maintain spore dormancy through the activation of the sporulation-specific adenylyl cyclase ACG (8).

Ammonia is produced by protein catabolism, and ammonia levels rise during development, when most energy is generated by the degradation of protein and RNA (27, 60, 71, 76). It has been suggested that glutamine synthetases, which incorporate ammonia into glutamine, control intracellular levels of ammonia. The expression of glutamine synthetases is developmentally regulated, and their activity becomes elevated during the culmination stage (12–14, 24), suggesting a role of glutamine

\* Corresponding author. Mailing address: Department of Cell Biology, Johns Hopkins University School of Medicine, 725 N. Wolfe St., Hunterian 107, Baltimore, MD 21205. Phone: (410) 502-6836. Fax: (410) 502-7826. E-mail for Ryuji Yoshino: ryoshino@jhmi.edu. E-mail for Miho Iijima: miiijima@jhmi.edu.

† Present address: University of Dundee, MSI/WTB Complex, Dow Street, Dundee DD1 5EH, United Kingdom.

<sup>∇</sup> Published ahead of print on 19 October 2007.

synthetases in culmination. Indeed, the pharmacological inhibition of glutamine synthetase blocks culmination during development (14). In addition, there are five genes, *amtA*, *amtB*, *amtC*, *rhgA*, and *rhgB*, which belong to the evolutionarily conserved family of ammonium transporter/methylammonium permease/rhesus protein (Amt/Mep/Rh) in the *Dictyostelium* genome (15). Previous studies have shown that developmental phenotypes in cells lacking AmtC can be rescued by deleting the AmtA protein (62). These studies suggest that AmtC and AmtA antagonistically regulate developmental processes and that AmtA and AmtC function in either ammonium transport or ammonium sensing (20, 37, 62). In this study, we show that AmtA regulates intracellular ammonium/ammonia levels during growth and development and is critical for the morphogenesis of multicellular aggregates and normal spore formation.

## MATERIALS AND METHODS

**Strains, culture, and development.** The *D. discoideum* strain Ax2 (40, 73) was used for wild-type cells in this study. For sequencing, strain Ax4 (40) was used. Cells were cultured at 22°C either in HL-5 medium (73) or on A medium agar plates (0.5% glucose, 0.05% yeast extract, 0.75 g of proteose peptone, 16.5 mM  $\text{KH}_2\text{PO}_4$ , 4.0 mM  $\text{K}_2\text{HPO}_4$ , 2.0 mM  $\text{MgSO}_4$ , 1.5% agar) associated with *Klebsiella pneumoniae*. For the selection and maintenance of transformants, the medium was supplemented with 10  $\mu\text{g}/\text{ml}$  of blasticidin S (Wako) or 20  $\mu\text{g}/\text{ml}$  Geneticin (Calbiochem). Peptone and yeast extract were obtained from Difco Laboratories.

For synchronous development, exponentially growing cells were harvested and washed three times in lower pad solution (LPS) buffer (20 mM KCl, 0.24 mM  $\text{MgCl}_2$ , 40 mM  $\text{K}_2\text{HPO}_4$ - $\text{KH}_2\text{PO}_4$ , pH 6.4) and placed on 47-mm-diameter cellulose membrane filters (Toyo Roshi Kaisha, Ltd.) at a density of  $5 \times 10^6$  cells/ $\text{cm}^2$ . The filters were placed on M-085 paper pads (Toyo Roshi Kaisha, Ltd.) containing LPS and housed in 6-cm-diameter petri dishes at 22°C. When  $\text{NH}_4\text{Cl}$  was added, ammonia buffer A (20 mM KCl, 2.5 mM  $\text{MgCl}_2$ , 40 mM  $\text{KH}_2\text{PO}_4$ , pH 7.3) was used instead of LPS buffer. Developmental processes were observed with an Olympus SZX12 microscope. Stalk lengths were determined using the NIH Image program (developed at the National Institutes of Health [http://rsb.info.nih.gov/nih-image/]).

**Nucleic acid analyses.** Genomic DNA was extracted as described previously (33, 35). Total RNA was isolated using TRIzol reagent (Invitrogen) by following the manufacturer's instructions. Poly(A)-containing RNA was purified using Oligotex-dT30 Super (JSR). RNA from prespore and prestalk cells was extracted as previously described (53, 56). Southern and Northern blot analyses were performed as previously described (57). A probe specific for *amtA* was prepared by random priming for a 1.1-kb EcoT22I/NotI-digested fragment of cDNA SSB490.

The *amtA* DNA sequence was determined by sequencing cDNA clones SSB490, SSB126, and SLB420, generated by the Japanese cDNA project (51, 67, 68), and the corresponding region of genomic DNA.

**Phylogenetic analysis.** A neighbor-joining tree was generated by the Clustal program (65) based on Pfam database alignment (PF00909). The Swiss-Prot/TrEMBL sequence accession numbers for the indicated proteins are as follows: *D. discoideum* AmtA, Q9BLG4; *D. discoideum* AmtB, Q9BLG3; *D. discoideum* AmtC, Q8MXYO; *Methanobacterium thermoautotrophicum* MTH661, O26757; *Escherichia coli* AmtB, P37905; *Saccharomyces cerevisiae* Mep1p, P40260; *S. cerevisiae* Mep2p, P41948; *S. cerevisiae* Mep3p, P53390; *Caenorhabditis elegans* C05E11.4, P54145; *Arabidopsis thaliana* AMT1;1, P54144; *A. thaliana* AMT1;2, Q9ZPJ8; *A. thaliana* AMT2;1, Q9M6N7; *Chlamydomonas reinhardtii* Rhp, Q94CJ2; *C. elegans* Rhp-1, Q9N2M5; *Drosophila melanogaster* Rhp, Q9V3T3; *Xenopus laevis* RhAG, Q9DGD8; *Homo sapiens* RhAG, O43514; *H. sapiens* RhBG, Q9H310; *H. sapiens* RhCG, Q9UBD6; *H. sapiens* RhCE, P18577; and *H. sapiens* RhD, Q02161. Amt alpha and beta groupings were based on previous work (39).

**Gene disruption and expression.** For *amtA* disruption, the blasticidin resistance gene (*bsr*) from pBsr $\Delta$ Bam (1) was used as a selection marker. The *amtA* coding region in the cDNA clone SSB490 was amplified by PCR using primers LA-XbaI (5'-gctctagaTGTAACCAACCACCACCCCAAATCC-3') and LA-HindIII (5'-ccaagcttGGTTTGGTTTAAATGCAGGTAGTGC-3') (lowercase indicates restriction sites), and then the PCR product was ligated with a 1.3-kb XbaI/HindIII fragment of pBsr $\Delta$ Bam. For efficient homologous recombination,

part of the 5' region of the construct was replaced by a genomic sequence containing the first intron. The sequence for displacement was amplified from genomic DNA with the primers LAP-1 (5'-AATCAAGTAGCACCAGATCCA GG-3') and LA-XbaI. The *bsr* gene was oriented in the same direction as the *amtA* gene to ensure the presence of a transcriptional terminator in the middle of the *amtA* coding sequence. From the resulting vector, pAMTAbSr (5.9 kb), the 2.9-kb *amtA* or *bsr* fragment was excised with SalI and NotI and used for transformation.

To express the wild-type *amtA* gene, a plasmid containing *amtA* was constructed by inserting a PCR fragment containing the complete *amtA* open reading frame into the *Dictyostelium* extrachromosomal expression vector HK12neo, a derivative of MB12neo (43).

**Determinations of ammonia/ammonium.** The intercellular levels of ammonia were measured as follows (2, 8). Cells were developed on nitrocellulose filters at  $5 \times 10^6$  cells/ $\text{cm}^2$ . Filters were placed on a paper pad containing 2 ml of KK2 buffer (20 mM  $\text{KH}_2\text{PO}_4$ - $\text{K}_2\text{HPO}_4$ , pH 6.4) and housed in petri dishes at 22°C. Cells were then harvested in 30 ml of ice-cold KK2 containing 20 mM EDTA at different time points during development, washed, and resuspended in 1 ml of KK2 buffer. The cell suspension was homogenized with a Mini-Bead-Beater (Wakenyaku Co., Ltd.), using 0.5-mm-diameter glass beads at 5,000 rpm for 3 min. For extracellular measurements, the paper pad under the nitrocellulose filter was soaked in 20 ml of KK2 buffer for 20 min to allow the ammonia to diffuse from the pad, and the resultant solution was used as a sample of the extracellular environment of the cells. To measure ammonia in the spores and spore mass solution, cells were developed on membrane filters, and spores were collected using a glass capillary. After centrifugation at  $1,400 \times g$  for 5 min, the supernatant was used as spore mass solution. The pellet of spores was resuspended in 1 ml of KK2 and then homogenized as described above. The concentrations of ammonia/ammonium in the sample solutions were measured using a Ti-9001Ka ammonia electrode (Toko Chemical Laboratories Co., Ltd.), by following the manufacturer's instructions.

**Spore viability and germination.** The germination rates of spores were determined as described previously (50). Spores were collected in KK2 buffer from fruiting bodies on agar plates with a wire loop. To induce synchronous germination, 30  $\mu\text{l}$  of supernatant from an overnight culture of *K. pneumoniae* in A medium was added to 0.3 ml of spore suspension in a siliconized 20-ml test tube (28). After being shaken at 150 rpm at 22°C for 6 h, germinated spores were counted using a hemocytometer. To determine the viability of spores, vital staining with propidium iodide (PI) was used (50). Spores were collected at 2, 4, 6, and 8 days after fruiting body formation, suspended in KK2 buffer, and stained with 20  $\mu\text{g}/\text{ml}$  of PI (Sigma) for 5 min.

**Glucose assay.** Glucose levels were measured as described previously (34), with a minor modification. Cells were grown in shaking cultures at 22°C in HL-5 medium with maltose as a carbohydrate source. Cells were harvested, developed for 0, 6, and 12 h in shaking cultures, and collected by centrifugation. The pellets were lysed by freezing at  $-70^\circ\text{C}$  overnight. The pellets were thawed and resuspended in distilled water at  $6 \times 10^8$  cells/ml. The lysates were clarified by centrifugation at  $12,000 \times g$  for 5 min, and 80  $\mu\text{l}$  of the supernatants was used to measure glucose using a Sigma glucose assay kit (catalog no. GAGO20), by following the manufacturer's instructions. The protein concentration of lysate was determined using the Bio-Rad protein assay kit (catalog no. 500-0006).

**Immunoblotting.** Cells were developed on agar plates, and spores were harvested. Whole-cell lysates were subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis and blotted on polyvinylidene difluoride membranes. Blots were immunostained with monoclonal antiactin antibody (C4; Chemicon) or monoclonal antiphosphotyrosine antibody clone PY20 (ICN), as previously described (81).

## RESULTS

**AmtA controls ammonia homeostasis during growth and development.** The *Dictyostelium* genome contains five genes, *amtA*, *amtB*, *amtC*, *rhgA*, and *rhgB*, which belong to the Amt/Mep/Rh family. A phylogenetic analysis showed that these five genes are divided into three groups: Amt alpha, Amt beta, and Rh (Fig. 1A). Although most organisms carry only one type of Amt/Mep/Rh gene (for example, vertebrates carry only genes that belong to the Rh group), *Dictyostelium* has all three types. In *Dictyostelium*, there are two genes that belong to each of the Rh (*rhgA* and *rhgB*) and Amt beta (*amtB* and *amtC*) groups, and




FIG. 1. Disruption of the *amtA* gene. (A) Phylogenetic analysis of ammonium transporters in *D. discoideum* (Dd), *M. thermautotrophicum* (Mt), *E. coli* (Ec), *S. cerevisiae* (Sc), *C. reinhardtii* (Cr), *C. elegans* (Ce), *D. melanogaster* (Dm), *X. laevis* (Xl), *H. sapiens* (Hs), and *A. thaliana* (At). The scale indicates the number of substitutions per site. (B) Disruption of *amtA*. A blasticidin resistance marker (*bsr* cassette) replaced the second intron of the *amtA* gene by homologous recombination. Exons are indicated by white boxes. B, Hc, and V indicate BamHI, HincII, and EcoRV restriction sites, respectively. (C) The *amtA* disruption was confirmed by Southern blotting, using the probe shown in panel B. Genomic DNA isolated from wild-type and *amtA*<sup>-</sup> cells was analyzed after digestion with HincII or BamHI and EcoRV (B+V).

the Amt alpha group contains a single gene, *amtA*. In this paper, we focused on the characterization of *amtA*. *amtA* encodes a 463-amino-acid protein (49.1 kDa), and a hydropathy analysis suggests that AmtA contains 11 transmembrane segments. The AmtA protein displays 11.0 to 23.6% identity to other Amt/Mep/Rh proteins in *Dictyostelium* and 23.8 to 35.6% identity to proteins in the Amt alpha groups in other organisms.

To determine whether AmtA regulates ammonium homeostasis, we disrupted the *amtA* gene in *Dictyostelium* by homologous recombination. The disruption was confirmed by Southern blot analysis (Fig. 1B and C). Wild-type cells showed 5.1- and 3.5-kb DNA fragments generated by HincII digestion and by BamHI/EcoRV digestion, respectively. On the other hand, 6.2- and 4.6-kb fragments were observed in cells disrupted for the *amtA* gene. We measured levels of intracellular and extracellular ammonia/ammonium during growth and development using ammonia-specific electrodes (2, 8). We observed  $5.24 \pm 0.14$  fmol of ammonia/ammonium per cell in wild-type cells during growth (Fig. 2A). In contrast, in *amtA*<sup>-</sup> cells, intracellular ammonium/ammonia levels were significantly increased. The mutants contained  $8.0 \pm 0.7$  fmol of ammonium/ammonia per cell, showing an ~1.5-fold increase. When we induced development by starvation, intracellular levels of ammonium/ammonia gradually de-





FIG. 2. AmtA is required for intracellular ammonia homeostasis. (A) Intracellular ammonia/ammonium levels. Cells were collected at 0, 12, and 20 h of development and homogenized, and the amounts of ammonia/ammonium were determined by an ammonia electrode. Values indicate means  $\pm$  standard errors of the means (SEM) ( $n = 3$ ). (B) Extracellular ammonia levels. Media were collected at the indicated time points, and the amounts of ammonia were determined as described in Materials and Methods. Values indicate means  $\pm$  SEM ( $n = 3$ ).

creased in wild-type cells during development. Although *amtA*<sup>-</sup> cells also showed a gradual decrease in ammonia/ammonium levels, intracellular levels of ammonia/ammonium in *amtA*<sup>-</sup> cells were much higher than those seen in wild-type cells at both the aggregation stage (12 h after starvation) and culmination stage (20 h). Increases in intracellular ammonia/ammonium levels could result from either overproduction of ammonia/ammonium or defects in its transport out of cells. Supporting the latter possibility, we found that extracellular levels of ammonium/ammonia were lower in *amtA*<sup>-</sup> cells (Fig. 2B). We confirmed that these phenotypes were caused by the loss of the *amtA* gene by expressing wild-type *amtA* in *amtA*<sup>-</sup> cells. All the phenotypes were significantly rescued by the expression of AmtA. These results clearly demonstrate that AmtA is required for normal ammonia homeostasis during growth and development.

Consistent with a role for AmtA in ammonia homeostasis, we found that *amtA* mRNA (1.7 kb) is expressed during both growth and development using Northern blot analysis (Fig. 3A). The expression level increased continuously during development. This is in contrast with previous studies using reverse transcription-PCR that showed that *amtA* mRNA levels do not change during development (20). The apparent difference may result from different methods used to measure mRNA levels. Nevertheless, the previous study and our current study clearly demonstrate that AmtA is expressed during development. Furthermore, we confirmed the localization of *amtA* mRNA in prespore cells in slugs (20), using Northern blotting (Fig. 3B).

We confirmed the results of previous studies (62) showing that *amtA*<sup>-</sup> cells normally grow both in shaking culture and on bacterial plates and that the time courses of development in wild-type and *amtA*<sup>-</sup> cells are similar, differentiating into fruiting bodies in 24 h upon starvation (data not shown). Wild-type and *amtA*<sup>-</sup> cells show similar sensitivities to ammonia in culmination (Fig. 3C). In Fig. 3C, cells were developed in the absence of NH<sub>4</sub>Cl, and then different amounts of NH<sub>4</sub>Cl were added to pads for further development. In addition, *amtA*<sup>-</sup> cells produced more aggregates and fruiting bodies than wild-








FIG. 4. Effects of ammonia on tip formation, chemotaxis, and culmination. (A) Effects of ammonia on aggregation and tip formation. Wild-type and *amtA*<sup>-</sup> cells were placed on nitrocellulose filters and allowed to develop for 36 h in the presence of 0, 30, and 50 mM NH<sub>4</sub>Cl. Bar, 0.5 mm. (B) Effects of ammonia on chemotaxis toward cAMP. Wild-type and *amtA*<sup>-</sup> cells were plated on nonnutrient plates in the presence or absence of 10 μM cAMP and observed after 6 h. As indicated, plates contain 0 or 30 mM NH<sub>4</sub>Cl. We determined the original position of the edge immediately after the cells were spotted. Bar, 0.2 mm.

type cells (Fig. 3D to F). In wild-type cells, the average density of aggregates was 593 aggregates/cm<sup>2</sup>. In contrast, *amtA*<sup>-</sup> cells produced 1,128 aggregates/cm<sup>2</sup>. The resulting fruiting bodies of *amtA*<sup>-</sup> cells were smaller than those of wild-type cells. The average stalk length in *amtA* mutants (0.77 mm) was almost half that in the wild-type (1.63 mm). The expression of wild-type AmtA in *amtA*<sup>-</sup> cells significantly suppressed those *amtA*<sup>-</sup> phenotypes. These results are consistent with the previous observation that AmtA plays a critical role in the number and size of aggregates and fruiting bodies (62). We also found that *amtA*<sup>-</sup> cells show smaller aggregation territory sizes and frequent group breakup (Fig. 3G). Since glucose has been suggested to increase aggregation size, we measured intracellular levels of glucose (34, 22). We found that glucose levels in *amtA*<sup>-</sup> cells were higher than those in wild-type cells (Fig.

3H). In contrast, protein amounts in wild-type and *amtA*<sup>-</sup> cells were indistinguishable (data not shown). These results suggest that the decreases of aggregation sizes in *amtA*<sup>-</sup> cells are affected mainly by ammonia, not glucose.

***amtA*<sup>-</sup> cells are hypersensitive to ammonia in early morphogenesis.** It has been shown that NH<sub>4</sub>Cl inhibits multiple steps in early developmental processes, including the aggregation and tip formation of aggregates (10, 23, 60). Since *amtA*<sup>-</sup> cells contain higher levels of ammonia/ammonium in early development than wild-type cells, we reasoned that mutant cells may be more sensitive to ammonia than wild-type cells. To test this idea, we induced development in *amtA*<sup>-</sup> and wild-type cells in the presence of ammonia. Cells were incubated throughout development on filter membranes containing 0, 30, or 50 mM of NH<sub>4</sub>Cl. As shown in Fig. 4A, in wild-type cells,

FIG. 3. *amtA*<sup>-</sup> cells display the aberrant morphology of fruiting bodies. (A) Poly(A)-containing RNAs were collected at the indicated time points and analyzed by Northern blotting with *amtA* cDNA as a probe. Five micrograms of poly(A)-containing RNAs were loaded in each lane. (B) *amtA* mRNA accumulates in prespore cells. Expression in respective cell types at the slug stage. Total RNA was extracted from prespore (psp) and prestalk (pst) cells in slugs. *ecmA* was used as a marker for prespore cells. (C) Effects of ammonia on culmination. Cells were developed in the absence of ammonia for 14 h and then transferred onto paper pads containing the indicated concentrations of NH<sub>4</sub>Cl. Photographs were taken 10 h after transfer. Bar, 1 mm. (D) *amtA*<sup>-</sup> cells produce many small fruiting bodies. Wild-type cells (WT), *amtA*<sup>-</sup> cells, and *amtA*<sup>-</sup> cells expressing *amtA* were plated on nitrocellulose filters to induce development and observed at 24 h. Bar, 0.5 mm. (E) Quantitation of aggregate density. Cells were plated on filters at a density of 5 × 10<sup>6</sup>/cm<sup>2</sup>. After 12 h, aggregates in an area of 81 mm<sup>2</sup> were counted. Values indicate means ± standard errors of the means (SEM) (*n* = 3). (F) Quantitation of stalk length. Values indicate means ± SEM (*n* ≥ 50). (G) Wild-type and *amtA*<sup>-</sup> cells were developed on nonnutrient plates for the indicated times. Compared to wild-type cells, *amtA*<sup>-</sup> cells show smaller aggregation territory sizes and increased group breakup. Cells were plated at densities of 5 × 10<sup>5</sup>/cm<sup>2</sup> and 5 × 10<sup>6</sup>/cm<sup>2</sup>. Bar, 1 mm. (H) *amtA*<sup>-</sup> cells show higher glucose levels than wild-type cells. Cells were developed for 0, 6, and 12 h and analyzed for glucose levels. Values indicate means ± SEM (*n* ≥ 5).

ammonia inhibited aggregation and tip formation in a concentration-dependent manner. In the absence of  $\text{NH}_4\text{Cl}$ , wild-type cells developed normally and formed fruiting bodies in 24 h after starvation. However, in the presence of 30 mM  $\text{NH}_4\text{Cl}$ , wild-type cells were able to form only aggregates with pointed ends (tip formation) and did not differentiate further into fruiting bodies. In 50 mM  $\text{NH}_4\text{Cl}$ , wild-type cells aggregated without tip formation. This result is not simply due to a delay in development. When we examined cells for prolonged periods of time (3 days), the cells still did not form fruiting bodies. In contrast, the mutants did not show tip formation of aggregates in 30 mM  $\text{NH}_4\text{Cl}$ , although *amtA*<sup>-</sup> cells formed fruiting bodies in the absence of  $\text{NH}_4\text{Cl}$ . Furthermore, 50 mM  $\text{NH}_4\text{Cl}$  completely blocked aggregation in *amtA*<sup>-</sup> cells. These results show that the loss of AmtA makes cells hypersensitive to ammonia in aggregation and tip formation during development.

***amtA*<sup>-</sup> cells are defective in chemotaxis in the presence of ammonia.** The inhibitory effect of ammonia on the aggregation of *amtA*<sup>-</sup> cells suggests that ammonia suppresses chemotaxis toward cAMP in mutant cells. To test this possibility, we examined chemotaxis toward cAMP using a spot assay (Fig. 4B). Cells were spotted on the center of plates which contained 10  $\mu\text{M}$  cAMP in the presence or absence of  $\text{NH}_4\text{Cl}$ . Since there were no nutrients available on the plate, cells initiated development processes immediately after being spotted. Cells secrete phosphodiesterase and degrade extracellular cAMP, generating a difference in cAMP concentrations around the spot, with higher concentrations outside and lower concentrations inside. Along the cAMP gradient, cells move outward from the spotted area. If cells are defective in chemotaxis, they will stay inside the spotted areas. When wild-type and *amtA*<sup>-</sup> cells were spotted on plates containing 10  $\mu\text{M}$  cAMP in the absence of  $\text{NH}_4\text{Cl}$ , both wild-type and mutant cells developed normally. At the periphery of the spotted area, we found that many cells moved outward (Fig. 4B). This outward movement is chemotactic migration toward cAMP, since cells move out only when cAMP is present. On plates lacking cAMP, cells developed normally, but they stayed in the spotted area and did not move outward. However, when we added 30 mM  $\text{NH}_4\text{Cl}$  to the plates, wild-type and *amtA*<sup>-</sup> cells showed distinct chemotactic behaviors. In the presence of 30 mM  $\text{NH}_4\text{Cl}$ , wild-type cells were still able to develop, form aggregates, and move outward toward cAMP. In contrast, when *amtA*<sup>-</sup> cells were spotted, the mutant cells developed normally and formed aggregates, but they failed to move out of the spotted area. Our data suggest that cells lacking AmtA become more sensitive to  $\text{NH}_4\text{Cl}$  during chemotaxis toward cAMP.

**AmtA is required for normal spore formation.** In *Dictyostelium*, the spore formation process is affected by ammonia through the regulation of cAMP-dependent PKA (30, 80). In addition, extra spore solution contains high concentrations of ammonium phosphate, which has been suggested to maintain the dormancy of spores (8, 9, 70). To determine whether AmtA is involved in ammonium homeostasis in spore and extra spore solution in sori, we collected sori, separated spores from extra spore solution by centrifugation, and measured levels of ammonia/ammonium. As shown in Fig. 5A, the levels of ammonia/ammonium increased twofold in *amtA*<sup>-</sup> spores. Wild-type spores contained 1.50 fmol of ammonia/ammonium per cell, whereas *amtA*<sup>-</sup> spores contained 2.69 fmol of ammonia/am-

monium per cell. In addition, ammonia levels in extra spore solution decreased twofold in the mutant cells (Fig. 5B). Therefore, our data indicate that AmtA is required for normal ammonia homeostasis in spores.

To probe the role of AmtA in spores, we examined the morphology, viability, and germination rates of spores. We observed spore morphology using phase-contrast microscopy and checked spore viability using PI, which stains only dead spores (7, 50, 52). We found that the viability of spores lacking AmtA is compromised as the spores age. Immediately after the completion of fruiting body formation, both wild-type and mutant spores showed oval shapes and very few spores were stained by PI. However, as spores aged, *amtA*<sup>-</sup> spores appeared to be darker under the phase-contrast microscope (Fig. 5C and D). The majority of the dark spores were stained by PI, indicating that they were dead. Twelve days after fruiting body formation, 79.0% of *amtA* mutant spores were stained by PI, whereas only 4.4% of wild-type spores were PI positive. Furthermore, we found that germination rates were decreased in *amtA*<sup>-</sup> spores (Fig. 5E). Probably as a result of their compromised viability, *amtA*<sup>-</sup> spores are also defective in normal germination. Immediately after spore formation, ~95% of wild-type spores germinated, whereas only 67% of *amtA* mutant spores germinated. Eight days after the completion of fruiting body formation, wild-type cells maintained a germination rate similar to that seen at day 0. In contrast, the germination rate of *amtA*<sup>-</sup> spores decreased as they aged. We found that only 20% of *amtA*<sup>-</sup> spores had germinated at 8 days. It has been shown that actins are organized into thick bundles in spores and are phosphorylated on tyrosine residues during the formation and maturation of spores (38, 59). Supporting the idea that *amtA*<sup>-</sup> cells are defective in normal spore formation, we found that tyrosine-phosphorylated actin levels decreased in *amtA*<sup>-</sup> spores (Fig. 5F). Therefore, our data indicate that AmtA is required for the formation and maintenance of viable spores.

## DISCUSSION

A large number of studies have shown that ammonia regulates multiple processes during the development of *Dictyostelium* (45, 47, 78–80). However, the molecular mechanisms underlying ammonia homeostasis are poorly understood. Previous studies have suggested that AmtA functions in ammonia transport or ammonia sensing (20, 62). In this paper, supporting a role for AmtA in ammonium transport, we have shown that AmtA is required for normal ammonia homeostasis in *Dictyostelium* during growth and development. Our data strongly suggest that AmtA is involved in the excretion of ammonia/ammonium from cells. Supporting our conclusion, cells lacking AmtA accumulate ammonia/ammonium. The impairment of ammonium efflux results in a reduction in extracellular levels of ammonia/ammonium. Furthermore, intraspore levels of ammonia/ammonium are also highly increased in *amtA* mutants. Since the extracellular levels of ammonia/ammonium are slightly higher than the intracellular levels in *Dictyostelium*, AmtA might actively transport ammonium out of cells. Although our data suggest that AmtA is an important ammonium transporter in development, they do not rule out






FIG. 5. AmtA is required for ammonia homeostasis, viability, and the germination of spores. Ammonia levels in spores (A) and spore mass solution (B). Three days after fruiting body formation, spore masses were collected and spores and sorus media were separated by centrifugation. The ammonia levels of each fraction were determined by an ammonia electrode. Values are normalized to  $10^8$  spores in  $10 \mu\text{l}$  sori. (C) Spores were collected 0 and 12 days after fruiting body formation was completed and stained with PI. Spores were observed by phase-contrast and fluorescence microscopy. Bar,  $50 \mu\text{m}$ . (D) Quantitation of PI-positive spores. Spores were collected at the indicated periods of time after fruiting body formation and stained with PI. Spores that were stained with PI were counted. (E) Spores were collected at the indicated time points after the completion of fruiting body formation and examined for germination, as described in Materials and Methods. (F) Changes of tyrosine phosphorylation of actin with aging. Total proteins were prepared from 1-, 2-, 6-, and 10-day-old spores and immunostained with antiactin antibody and antiphosphotyrosine antibody.

the possibility that AmtA also functions in ammonia sensing, as previously proposed (62).

Previous studies have shown that AmtA is important for the morphogenesis of fruiting bodies and for resistance to high concentrations of ammonia in culmination (62). Our study confirmed the previous findings and further identifies three additional developmental processes which involve AmtA-me-

diated ammonia homeostasis, including chemotaxis, tip formation, and spore formation. At the beginning of development, single cells undergo chemotaxis toward aggregation centers, which release the chemoattractant cAMP, leading to the formation of multicellular aggregates. We found that AmtA affects the sensitivity to ammonia in inhibition of chemotactic migration toward cAMP. *amtA*<sup>-</sup> cells fail to undergo chemo-

taxis in the presence of  $\text{NH}_4\text{Cl}$ . Since the concentrations of  $\text{NH}_4\text{Cl}$  that block chemotaxis do not inhibit the normal development of  $\text{amtA}^-$  cells, it is unlikely that the chemotaxis phenotypes result simply from developmental defects. In addition, in our experiments, we examined chemotaxis toward exogenously added cAMP. Therefore, the observed chemotaxis defect is not due to the inhibition of the production and secretion of cAMP by ammonia. Rather, we suggest that ammonia directly affects signaling pathways for chemotaxis. Supporting our hypothesis, it has been shown that ammonia affects intracellular pH and thereby inhibits chemotaxis (10, 19, 69). It is possible that intracellular pH is regulated at least partially by AmtA during chemotaxis. Since the morphogenesis of aggregates involves cAMP signaling (66), the impaired chemotaxis may result in defects in tip formation in aggregates in  $\text{amtA}^-$  cells. Furthermore, at the final stage of development,  $\text{amtA}^-$  cells formed much smaller fruiting bodies than those of wild-type cells, even in the absence of exogenously added ammonia. Many smaller fruiting bodies may result from incomplete chemotaxis, which could produce many smaller aggregates. Alternatively, it is also possible that ammonia homeostasis directly controls the size of the fruiting bodies by affecting the differentiation of spore and stalk cells.

One of our most interesting observations is that AmtA regulates ammonia levels in spores. We found that  $\text{amtA}^-$  spores in fruiting bodies contained higher levels of ammonia/ammonium but that extracellular levels were decreased. In addition,  $\text{amtA}^-$  spores were severely defective in viability, and these phenotypes became more severe as spores aged. It has been shown that ammonia in sori is critical for the formation and maintenance of spores (8, 9, 30, 70, 80). Thus, our studies demonstrate that AmtA is a critical regulator for ammonia homeostasis during spore formation. It is likely that the reduced viability of  $\text{amtA}^-$  spores leads to their defect in germination. Unlike with cells lacking the histidine kinase *dhkB*, which are defective in spore dormancy and prematurely germinate in sorus (82), we did not observe premature germination in  $\text{amtA}^-$  spores; it is unlikely that  $\text{amtA}^-$  cells are unable to inhibit premature germination in sorus, but rather they seem to be defective in formation of fully resistant spores. It has been suggested that the organization of the actin cytoskeleton is important for the formation and stabilization of spores (38, 58, 59). During spore formation, a large fraction of actin molecules are phosphorylated on tyrosine residues and organized into thick bundles. This process is induced under PKA activity and later mediated by the MADS box transcription factor SrfA (16–18). We found that the level of tyrosine phosphorylation of actin remains low in  $\text{amtA}^-$  spores, suggesting that ammonia homeostasis might regulate spore formation and stabilization by affecting the actin organization mediated by PKA and SrfA.

Our current study and previous study have shown that the disruption of AmtA does not cause the severe defects in the culmination step of aggregates on its own (62). Wild-type and  $\text{amtA}^-$  cells showed similar sensitivities to exogenously added ammonia in culmination, suggesting that other ammonium transporters regulate ammonia levels at the initiation of culmination. Supporting this idea, previous studies showed that AmtC was important for the induction of culmination and that cells lacking AmtC failed to culminate and remained as slugs (20, 37). In contrast to AmtA, AmtC is proposed to function as

a sensor which monitors ammonia levels and activates intracellular signaling. Consistent with this idea, AmtC is preferentially expressed at the tip region of slugs, where the culmination signal may be generated (20). Interestingly, the disruption of *amtC* in the *amtC* null strain restored the ability of cells to differentiate to undergo the culmination stage (62). In this study, we showed that *amtA* functions as an ammonium transporter, although it is possible that *amtA* controls the slug/culmination transition as an ammonia sensor. To determine whether AmtC functions as an ammonium sensor or transporter, it would be important to determine whether intracellular levels of ammonia are also altered in  $\text{amtC}^-$  cells.

Members of the Amt/Mep/Rh family are found in all domains of life and involved in a variety of biological processes. In microorganisms, ammonia is a nutrient, and most Amt/Mep/Rh proteins participate in the uptake of ammonium into cells (11, 31, 36, 42, 49). In contrast, in animals, ammonia is a waste product resulting from amino acid catabolism (29). Rh proteins have been suggested to excrete ammonia to maintain intracellular homeostasis (3, 4, 32, 48, 55, 75), although several recent studies suggest that the substrate of Rh proteins is  $\text{CO}_2$ , not ammonia (41, 54). Furthermore, yeast Mep2 ammonium permease has been shown to function as a sensor of ammonia, generating a signal to regulate pseudohyphal differentiation (21, 46). Mep2 may lack the ability to transport ammonium across membranes (46). In contrast to most organisms, which carry only one type of ammonium transporter among the three subfamilies (Amt alpha, Amt beta, and Rh groups), *Dictyostelium* contains five ammonium transporters that belong to all three subfamilies. It would be tempting to speculate that *Dictyostelium* uses the different functions of ammonium transporters, such as ammonium uptake, efflux, and sensing, to control its unique development.

#### ACKNOWLEDGMENTS

We thank Hideko Urushihara (University of Tsukuba) and Charles K. Singleton (Vanderbilt University) for valuable suggestions and comments. We also thank the *Dictyostelium* cDNA project in Japan for providing cDNA clones.

Research was supported by a Leukemia and Lymphoma Society fellowship (3374-05) and an American Heart Association beginning grant-in-aid (0765345U) (M.I.), an American Heart Association scientist development grant (0730247N) (H.S.), and Johns Hopkins University (M.I. and H.S.).

#### REFERENCES

- Adachi, H., T. Hasebe, K. Yoshinaga, T. Ohta, and K. Sutoh. 1994. Isolation of *Dictyostelium discoideum* cytokinesis mutants by restriction enzyme-mediated integration of the blasticidin S resistance marker. *Biochem. Biophys. Res. Commun.* **205**:1808–1814.
- Aeckerle, S., B. Wurster, and D. Malchow. 1985. Oscillations and cyclic AMP-induced changes of the  $\text{K}^+$  concentration in *Dictyostelium discoideum*. *EMBO J.* **4**:39–43.
- Bakouh, N., F. Benjelloun, B. Cherif-Zahar, and G. Planelles. 2006. The challenge of understanding ammonium homeostasis and the role of the Rh glycoproteins. *Transfus. Clin. Biol.* **13**:139–146.
- Biver, S., S. Scohy, J. Szpirer, C. Szpirer, B. Andre, and A. M. Marini. 2006. Physiological role of the putative ammonium transporter RhCG in the mouse. *Transfus. Clin. Biol.* **13**:167–168.
- Bloom, L., and R. R. Kay. 1988. The search for morphogens in *Dictyostelium*. *Bioessays* **9**:187–191.
- Bradbury, J. M., and J. D. Gross. 1989. The effect of ammonia on cell-type-specific enzyme accumulation in *Dictyostelium discoideum*. *Cell Differ. Dev.* **27**:121–128.
- Cornillon, S., C. Foa, J. Davoust, N. Buonavista, J. D. Gross, and P. Golstein. 1994. Programmed cell death in *Dictyostelium*. *J. Cell Sci.* **107**:2691–2704.



8. Cotter, D. A., A. J. Dunbar, S. D. Buconjic, and J. F. Wheldrake. 1999. Ammonium phosphate in sori of *Dictyostelium discoideum* promotes spore dormancy through stimulation of the osmosensor ACG. *Microbiology* **145**: 1891–1901.
9. Cotter, D. A., D. C. Mahadeo, D. N. Cervi, Y. Kishi, K. Gale, T. Sands, and M. Sameshima. 2000. Environmental regulation of pathways controlling sporulation, dormancy and germination utilizes bacterial-like signaling complexes in *Dictyostelium discoideum*. *Protist* **151**:111–126.
10. Davies, L., M. Satre, J. B. Martin, and J. D. Gross. 1993. The target of ammonia action in *Dictyostelium*. *Cell* **75**:321–327.
11. Detsch, C., and J. Stulke. 2003. Ammonium utilization in *Bacillus subtilis*: transport and regulatory functions of NrgA and NrgB. *Microbiology* **149**: 3289–3297.
12. Dunbar, A. J., and J. F. Wheldrake. 1995. Evidence for a developmentally regulated prespore-specific glutamine synthetase in the cellular slime mould *Dictyostelium discoideum*. *Microbiology* **141**:1125–1130.
13. Dunbar, A. J., and J. F. Wheldrake. 1997. Analysis of mRNA levels for developmentally regulated prespore specific glutamine synthetase in *Dictyostelium discoideum*. *Dev. Growth Differ.* **39**:617–624.
14. Dunbar, A. J., and J. F. Wheldrake. 1997. Effect of the glutamine synthetase inhibitor, methionine sulfoximine, on the growth and differentiation of *Dictyostelium discoideum*. *FEMS Microbiol. Lett.* **151**:163–168.
15. Eichinger, L., J. A. Pachebat, G. Glöckner, M. A. Rajandream, R. Sugang, M. Berriman, J. Song, R. Olsen, K. Szafarski, Q. Xu, B. Tunggal, S. Kummerfeld, M. Madera, B. A. Konfortov, F. Rivero, A. T. Bankier, R. Lehmann, N. Hamlin, R. Davies, P. Gaudet, P. Fey, K. Pilcher, G. Chen, D. Saunders, E. Sodergren, P. Davis, A. Kerhornou, X. Nie, N. Hall, C. Anjard, L. Hemphill, N. Bason, P. Farbrother, B. Desany, E. Just, T. Morio, R. Rost, C. Churcher, J. Cooper, S. Haydock, N. van Driessche, A. Cronin, I. Goodhead, D. Muzny, T. Mourier, A. Pain, M. Lu, D. Harper, R. Lindsay, H. Hauser, K. James, M. Quiles, M. Madan Babu, T. Saito, C. Buchrieser, A. Wardroper, M. Felder, M. Thangavelu, D. Johnson, A. Knights, H. Loulseged, K. Mungall, K. Oliver, C. Price, M. A. Quail, H. Urushihara, J. Hernandez, E. Rabinowitsch, D. Steffen, M. Sanders, J. Ma, Y. Kohara, S. Sharp, M. Simmonds, S. Spiegler, A. Tivey, S. Sugano, B. White, D. Walker, J. Woodward, T. Winckler, Y. Tanaka, G. Shaulsky, M. Schleicher, G. Weinstein, A. Rosenthal, E. C. Cox, R. L. Chisholm, R. Gibbs, W. F. Loomis, M. Platzer, R. R. Kay, J. Williams, P. H. Dear, A. A. Noegel, B. Barrell, and A. Kuspa. 2005. The genome of the social amoeba *Dictyostelium discoideum*. *Nature* **435**:43–57.
16. Escalante, R., and L. Sastre. 1998. A serum response factor homolog is required for spore differentiation in *Dictyostelium*. *Development* **125**:3801–3808.
17. Escalante, R., and L. Sastre. 2002. Regulated expression of the MADS-box transcription factor SrfA mediates activation of gene expression by protein kinase A during *Dictyostelium* sporulation. *Mech. Dev.* **117**:201–208.
18. Escalante, R., Y. Yamada, D. Cotter, L. Sastre, and M. Sameshima. 2004. The MADS-box transcription factor SrfA is required for actin cytoskeleton organization and spore coat stability during *Dictyostelium* sporulation. *Mech. Dev.* **121**:51–56.
19. Feit, I. N., E. J. Medynski, and M. J. Rothrock. 2001. Ammonia differentially suppresses the cAMP chemotaxis of anterior-like cells and prestalk cells in *Dictyostelium discoideum*. *J. Biosci.* **26**:157–166.
20. Follstaedt, S. C., J. H. Kirsten, and C. K. Singleton. 2003. Temporal and spatial expression of ammonium transporter genes during growth and development of *Dictyostelium discoideum*. *Differentiation* **71**:557–566.
21. Forsberg, H., and P. O. Ljungdahl. 2001. Sensors of extracellular nutrients in *Saccharomyces cerevisiae*. *Curr. Genet.* **40**:91–109.
22. Gao, T., C. Roisin-Bouffay, R. D. Hatton, L. Tang, D. A. Brock, T. Deshazo, L. Olson, W.-P. Hong, W. Jang, E. Canseco, D. Bakthavatsalam, and R. H. Gomer. 2007. A cell number-counting factor regulates levels of a novel protein, SsIA, as part of a group size regulation mechanism in *Dictyostelium*. *Eukaryot. Cell* **6**:1538–1551.
23. Gee, K., F. Russell, and J. D. Gross. 1994. Ammonia hypersensitivity of slugger mutants of *D. discoideum*. *J. Cell Sci.* **107**:701–708.
24. Gregg, J. H., A. L. Hackney, and J. O. Krivanek. 1954. Nitrogen metabolism of the slime mold *Dictyostelium discoideum* during growth and morphogenesis. *Biol. Bull.* **107**:226–235.
25. Gross, J. D. 1994. Developmental decisions in *Dictyostelium discoideum*. *Microbiol. Rev.* **58**:330–351.
26. Gross, J. D., C. D. Town, J. J. Brookman, K. A. Jermyn, M. J. Peacey, and R. R. Kay. 1981. Cell patterning in *Dictyostelium*. *Philos. Trans. R. Soc. Lond. B* **295**:497–508.
27. Hames, B. D., and J. M. Ashworth. 1974. The metabolism of macromolecules during the differentiation of myxamoebae of the cellular slime mould *Dictyostelium discoideum* containing different amounts of glycogen. *Biochem. J.* **142**:301–315.
28. Hashimoto, Y., Y. Tanaka, and T. Yamada. 1976. Spore germination promoter of *Dictyostelium discoideum* excreted by *Aerobacter aerogenes*. *J. Cell Sci.* **21**:261–271.
29. Heitman, J., and P. Agre. 2000. A new face of the Rhesus antigen. *Nat. Genet.* **26**:258–259.
30. Hopper, N. A., A. J. Harwood, S. Bouzid, M. Veron, and J. G. Williams. 1993. Activation of the prespore and spore cell pathway of *Dictyostelium* differentiation by cAMP-dependent protein kinase and evidence for its upstream regulation by ammonia. *EMBO J.* **12**:2459–2466.
31. Howitt, S. M., and M. K. Udvardi. 2000. Structure, function and regulation of ammonium transporters in plants. *Biochim. Biophys. Acta* **1465**:152–170.
32. Huang, C. H., and P. Z. Liu. 2001. New insights into the Rh superfamily of genes and proteins in erythroid cells and nonerythroid tissues. *Blood Cells Mol. Dis.* **27**:90–101.
33. Hughes, J. E., and D. L. Welker. 1988. A mini-screen technique for analyzing nuclear DNA from a single *Dictyostelium* colony. *Nucleic Acids Res.* **16**:2338.
34. Jang, W., B. Chiem, and R. H. Gomer. 2002. A secreted cell number counting factor represses intracellular glucose levels to regulate group size in *Dictyostelium*. *J. Biol. Chem.* **277**:39202–49208.
35. Kawabe, Y., T. Enomoto, T. Morio, H. Urushihara, and Y. Tanaka. 1999. LbrA, a protein predicted to have a role in vesicle trafficking, is necessary for normal morphogenesis in *Polysphondylium pallidum*. *Gene* **239**:75–79.
36. Khademi, S., and R. M. Stroud. 2006. The Amt/MEP/Rh family: structure of AmtB and the mechanism of ammonia gas conduction. *Physiology (Bethesda)* **21**:419–429.
37. Kirsten, J. H., Y. Xiong, A. J. Dunbar, M. Rai, and C. K. Singleton. 2005. Ammonium transporter C of *Dictyostelium discoideum* is required for correct prestalk gene expression and for regulating the choice between slug migration and culmination. *Dev. Biol.* **287**:146–156.
38. Kishi, Y., C. Clements, D. C. Mahadeo, D. A. Cotter, and M. Sameshima. 1998. High levels of actin tyrosine phosphorylation: correlation with the dormant state of *Dictyostelium* spores. *J. Cell Sci.* **111**:2923–2932.
39. Kitano, T., and N. Saitou. 2000. Evolutionary history of the Rh blood group-related genes in vertebrates. *Immunogenetics* **51**:856–862.
40. Knecht, D. A., S. M. Cohen, W. F. Loomis, and H. F. Lodish. 1986. Developmental regulation of *Dictyostelium discoideum* actin gene fusions carried on low-copy and high-copy transformation vectors. *Mol. Cell. Biol.* **6**:3973–3983.
41. Kustu, S., and W. Inwood. 2006. Biological gas channels for NH<sub>3</sub> and CO<sub>2</sub>: evidence that Rh (Rhesus) proteins are CO<sub>2</sub> channels. *Transfus. Clin. Biol.* **13**:103–110.
42. Li, X. D., D. Lupo, L. Zheng, and F. Winkler. 2006. Structural and functional insights into the AmtB/Mep/Rh protein family. *Transfus. Clin. Biol.* **13**: 65–69.
43. Linskens, M. H., P. D. Grootenhuys, M. Blaauw, B. Huisman-de Winkel, A. Van Ravestein, P. J. Van Haastert, and J. C. Heikoop. 1999. Random mutagenesis and screening of complex glycoproteins: expression of human gonadotropins in *Dictyostelium discoideum*. *FASEB J.* **13**:639–645.
44. Loomis, W. F. 1975. *Dictyostelium discoideum*: a developmental system. Academic Press, New York, NY.
45. Loomis, W. F. 1988. Signals that regulate differentiation in *Dictyostelium*, p.25–30. ISI atlas of science: immunology. Institute for Scientific Information, Philadelphia, PA.
46. Lorenz, M. C., and J. Heitman. 1998. The MEP2 ammonium permease regulates pseudohyphal differentiation in *Saccharomyces cerevisiae*. *EMBO J.* **17**:1236–1247.
47. Mahadeo, D. C., and C. A. Parent. 2006. Signal relay during the life cycle of *Dictyostelium*. *Curr. Top. Dev. Biol.* **73**:115–140.
48. Marini, A. M., G. Matassi, V. Raynal, B. Andre, J. P. Cartron, and B. Cherif-Zahar. 2000. The human Rhesus-associated RhAG protein and a kidney homologue promote ammonium transport in yeast. *Nat. Genet.* **26**: 341–344.
49. Marini, A.-M., S. Soussi-Boudekou, S. Vissers, and B. Andre. 1997. A family of ammonium transporters in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **17**: 4282–4293.
50. Mitra, B. N., R. Yoshino, T. Morio, M. Yokoyama, M. Maeda, H. Urushihara, and Y. Tanaka. 2000. Loss of a member of the aquaporin gene family, *aqpA* affects spore dormancy in *Dictyostelium*. *Gene* **251**:131–139.
51. Morio, T., H. Urushihara, T. Saito, Y. Ugawa, H. Mizuno, M. Yoshida, R. Yoshino, B. N. Mitra, M. Pi, T. Sato, K. Takemoto, H. Yasukawa, J. Williams, M. Maeda, I. Takeuchi, H. Ochiai, and Y. Tanaka. 1998. The *Dictyostelium* developmental cDNA project: generation and analysis of expressed sequence tags from the first-finger stage of development. *DNA Res.* **5**:335–340.
52. Nuckolls, G. H., N. Osherov, W. F. Loomis, and J. A. Spudich. 1996. The *Dictyostelium* dual-specificity kinase splA is essential for spore differentiation. *Development* **122**:3295–3305.
53. Ozaki, T., M. Hasegawa, Y. Hamada, M. Tasaka, M. Iwabuchi, and I. Takeuchi. 1988. Molecular cloning of cell-type-specific cDNAs exhibiting new types of developmental regulation in *Dictyostelium discoideum*. *Cell Differ.* **23**:119–124.
54. Peng, J., and C. H. Huang. 2006. Rh proteins vs Amt proteins: an organismal and phylogenetic perspective on CO<sub>2</sub> and NH<sub>3</sub> gas channels. *Transfus. Clin. Biol.* **13**:85–94.
55. Planelles, G. 2007. Ammonium homeostasis and human Rhesus glycoproteins. *Nephron Physiol.* **105**:11–17.
56. Ratner, D., and W. Borth. 1983. Comparison of differentiating *Dictyostelium*

- discoideum* cell types separated by an improved method of density gradient centrifugation. *Exp. Cell Res.* **143**:1–13.
57. Sambrook, J., and D. W. Russell. 2001. Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
  58. Sameshima, M., Y. Kishi, M. Osumi, D. Mahadeo, and D. A. Cotter. 2000. Novel actin cytoskeleton: actin tubules. *Cell Struct. Funct.* **25**:291–295.
  59. Sameshima, M., Y. Kishi, M. Osumi, R. Minamikawa-Tachino, D. Mahadeo, and D. A. Cotter. 2001. The formation of actin rods composed of actin tubules in *Dictyostelium discoideum* spores. *J. Struct. Biol.* **136**:7–19.
  60. Schindler, J., and M. Sussman. 1977. Ammonia determines the choice of morphogenetic pathways in *Dictyostelium discoideum*. *J. Mol. Biol.* **116**:161–169.
  61. Schindler, J., and M. Sussman. 1979. Inhibition by ammonia of intracellular cAMP accumulation in *Dictyostelium discoideum*: its significance for the regulation of morphogenesis. *Dev. Genet.* **1**:13–20.
  62. Singleton, C. K., J. H. Kirsten, and C. J. Dinsmore. 2006. Function of ammonium transporter A in the initiation of culmination of development in *Dictyostelium discoideum*. *Eukaryot. Cell* **5**:991–996.
  63. Singleton, C. K., M. J. Zinda, B. Mykytka, and P. Yang. 1998. The histidine kinase *dhkC* regulates the choice between migrating slugs and terminal differentiation in *Dictyostelium discoideum*. *Dev. Biol.* **203**:345–357.
  64. So, J. S., and G. Weeks. 1992. The effects of presumptive morphogens on prestalk and prespore cell gene expression in monolayers of *Dictyostelium discoideum*. *Differentiation* **51**:73–78.
  65. Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
  66. Traynor, D., R. H. Kessin, and J. G. Williams. 1992. Chemotactic sorting to cAMP in the multicellular stages of *Dictyostelium* development. *Proc. Natl. Acad. Sci. USA* **89**:8303–8307.
  67. Urushihara, H., T. Morio, T. Saito, Y. Kohara, E. Koriki, H. Ochiai, M. Maeda, J. G. Williams, I. Takeuchi, and Y. Tanaka. 2004. Analyses of cDNAs from growth and slug stages of *Dictyostelium discoideum*. *Nucleic Acids Res.* **32**:1647–1653.
  68. Urushihara, H., T. Morio, and Y. Tanaka. 2006. The cDNA sequencing project. *Methods Mol. Biol.* **346**:31–49.
  69. Van Duijn, B., and K. Inouye. 1991. Regulation of movement speed by intracellular pH during *Dictyostelium discoideum* chemotaxis. *Proc. Natl. Acad. Sci. USA* **88**:4951–4955.
  70. Viridy, K. J., T. W. Sands, S. H. Kopko, S. van Es, M. Meima, P. Schaap, and D. A. Cotter. 1999. High cAMP in spores of *Dictyostelium discoideum*: association with spore dormancy and inhibition of germination. *Microbiology* **145**:1883–1890.
  71. Walsh, J., and B. E. Wright. 1978. Kinetics of net RNA degradation during development in *Dictyostelium discoideum*. *J. Gen. Microbiol.* **108**:57–62.
  72. Wang, M., and P. Schaap. 1989. Ammonia depletion and DIF trigger stalk cell intact *Dictyostelium discoideum* slugs. *Development* **105**:569–574.
  73. Watts, D. J., and J. M. Ashworth. 1970. Growth of myxameobae of the cellular slime mould *Dictyostelium discoideum* in axenic culture. *Biochem. J.* **119**:171–174.
  74. Weeks, G., and J. D. Gross. 1991. Potential morphogens involved in pattern formation during *Dictyostelium* differentiation. *Biochem. Cell Biol.* **69**:608–617.
  75. Weiner, I. D. 2004. The Rh gene family and renal ammonium transport. *Curr. Opin. Nephrol. Hypertens.* **13**:533–540.
  76. White, G. J., and M. Sussman. 1961. Metabolism of major cell components during slime mold morphogenesis. *Biochim. Biophys. Acta* **53**:285–293.
  77. Williams, G. B., E. M. Elder, and M. Sussman. 1984. Modulation of the cAMP relay in *Dictyostelium discoideum* by ammonia and other metabolites: possible morphogenetic consequences. *Dev. Biol.* **105**:377–388.
  78. Williams, J. G. 1988. The role of diffusible molecules in regulating the cellular differentiation of *Dictyostelium discoideum*. *Development* **103**:1–16.
  79. Williams, J. G. 1991. Regulation of cellular differentiation during *Dictyostelium* morphogenesis. *Curr. Opin. Genet. Dev.* **1**:358–362.
  80. Williams, J. G., A. J. Harwood, N. A. Hopper, M. N. Simon, S. Bouzid, and M. Veron. 1993. Regulation of *Dictyostelium* morphogenesis by cAMP-dependent protein kinase. *Philos. Trans. R. Soc. Lond. B* **340**:305–313.
  81. Yamada, Y., and M. Sameshima. 2003. Hypertonic signal promotes stability of *Dictyostelium* spores via a PKA-independent pathway. *FEMS Microbiol. Lett.* **229**:159–164.
  82. Zinda, M. J., and C. K. Singleton. 1998. The hybrid histidine kinase *dhkB* regulates spore germination in *Dictyostelium discoideum*. *Dev. Biol.* **196**:171–183.