

1 **Nucleosome positioning and histone H3-acetylation are independent processes in**  
2 **the *Aspergillus nidulans prnD-prnB* bidirectional promoter.**

3

4 Yazmid Reyes-Dominguez<sup>1,2</sup>, Frank Narendja<sup>1,3</sup>, Harald Berger<sup>1</sup>, Andreas Gallmetzer<sup>1</sup>,  
5 Rafael Fernandez-Martin<sup>2,4</sup>, Irene Garcia<sup>2,5</sup>, Claudio Scazzocchio<sup>2,6</sup> and Joseph  
6 Strauss<sup>1,#</sup>

7 <sup>1</sup> Fungal Genomics Unit, Austrian Research Centers and BOKU Vienna, A-1190  
8 Vienna, Austria, <sup>2</sup> Istitute de Genetique et Microbiologie, Université Paris-Sud, F-91495  
9 Orsay-CEDEX, France.

10 <sup>3</sup> present address: Federal Environment Agency - Umweltbundesamt GesmbH,  
11 Spittelauerlände 5, A-1090 Vienna, Austria.

12 <sup>4</sup> present address: Laboratorio Biotecnología Animal, Universidad de Buenos Aires,  
13 Avenida San Martin 4453, Capital Federal, Argentina

14 <sup>5</sup> present address: Instituto de Bioquímica Vegetal y Fotosíntesis (IBVF), CSIC-  
15 Universidad de Sevilla, Américo Vespucio, 49, 41092-Sevilla, SPAIN

16 <sup>6</sup> present address: Department of Microbiology, Imperial College, London, Flowers  
17 Building, Armstrong Road, London SW7 2AZ UK

18

19

20 # corresponding author: Joseph Strauss, Fungal Genomics Unit, Austrian Research  
21 Centers and BOKU Vienna, Muthgasse 18, A-1190 Vienna, Austria. Tel.: +43-1-36006-  
22 6720; Fax: +43-1-36006-6392; E-mail: joseph.strauss@boku.ac.at

23

1

## 2 **Abstract**

3 In *Aspergillus nidulans*, proline can be used as carbon and nitrogen source and its  
4 metabolism requires the integration of three signals including proline induction, and  
5 nitrogen and carbon metabolite de-repression. We have previously shown that the  
6 bidirectional promoter in the *prnD-prnB* intergenic region undergoes drastic chromatin  
7 rearrangements where proline induction leads to loss of positioned nucleosomes  
8 whereas simultaneous carbon and nitrogen metabolite repression results in partial re-  
9 positioning of these nucleosomes. In the proline cluster, inhibition of deacetylases by  
10 trichostatin A (TSA), leads to partial de-repression and is associated with a lack of  
11 nucleosome positioning. Here, we investigate the effect of histone acetylation in the  
12 proline cluster using strains deleted for essential components of putative *A. nidulans*  
13 histone acetyl transferase (HAT) complexes, namely *gcnE* and *adaB*, the orthologues of  
14 the *S. cerevisiae* *GCN5* and *ADA2* genes, respectively. Surprisingly, GcnE and AdaB  
15 are not required for transcriptional activation and chromatin remodelling but for  
16 repression of *prnB* and *prnD* and for re-positioning of nucleosomes in the divergent  
17 promoter region. Chromatin immunoprecipitation (ChIP) directed against histone H3  
18 lysines K9 and K14 revealed that GcnE and AdaB participate in increasing the  
19 acetylation level of at least one nucleosome in the *prnD-prnB* intergenic region during  
20 activation, but these activities do not determine nucleosome positioning. Our results are  
21 consistent with a function of GcnE and AdaB in gene repression of the proline cluster,  
22 probably an indirect effect related to the function of CreA, the DNA-binding protein  
23 mediating carbon catabolite repression in *Aspergillus nidulans*.

24

## 1 **Introduction**

2 In eukaryotic cells, acetylation of histones is correlated with both transcriptional  
3 activation and the chromatin rearrangements usually associated with it. A turning point  
4 in our understanding of the mechanism of histone acetylation was the demonstration of  
5 the intrinsic acetylase activity of the *Tetrahymena pyriformis* p55 protein (7), the  
6 homologue of the *Saccharomyces cerevisiae* Gcn5p protein, the latter already known to  
7 be involved in the transcriptional activation of a number of genes in this organism  
8 (reviewed in (33)). Homologues, possibly orthologues, of Gcn5p, are universally present  
9 in eukaryotes. Gcn5p and its homologues interact with chromatin in large multi-protein  
10 complexes such as ADA and SAGA (17). SAGA is a multiprotein complex whose  
11 Gcn5p subunit possesses histone acetyltransferase activity. Gcn5p acetylates several lysine  
12 residues on histones N-termini including K9 and K14 on histone H3 and K8 and K16  
13 on histone H4 (21). The Gcn5p-containing complexes share several subunits, such as  
14 Ada2p, Ada3p, Spt3p and Tra1p (24) and it has been estimated that the yeast SAGA  
15 complex regulates the expression of ~ 10% of *S.cerevisiae* genes, approximately one-  
16 third of them being negatively regulated (23). Gcn5p forms a ternary complex with  
17 Ada2p and Ada3p, a complex which is conserved in SAGA and ADA (2). The proteins  
18 of the complex modulate the acetylating activity of Gcn5p (8, 17, 35) in patterns that are  
19 not identical for different promoters or different transcriptional activators. (9, 33). In the  
20 context of transcriptional regulation the SAGA complex can have different functions.  
21 The role of SAGA has been extensively studied in the *S. cerevisiae* *GAL1* promoter and  
22 has been shown to be essential for *GAL1* transcription. In the case of *GAL1*, the first  
23 step in transcriptional activation is binding of the specific Gal4p activator which in turn  
24 recruits SAGA to the upstream activating sequence (UAS) and the UAS-bound SAGA

1 then promotes the binding of TATA-binding protein and assembly of pre-initiation  
2 complex (5, 22). In contrast to the *GAL1* system, in the yeast *ADE* regulon SAGA is not  
3 recruited by the specific activators Bas1p and Pho2p (nor is SWI/SNF, another  
4 chromatin remodelling complex recruited in this promoter). Instead, in promoters of this  
5 regulon SAGA function is required for recruitment and efficient binding of the specific  
6 activators (19). In the yeast nitrogen-carbon utilization regulatory interface, different  
7 components of SAGA have been shown to have different functions. The expression of  
8 both paralogous glutamate dehydrogenase genes, *GDH1* and *GDH3*, require different  
9 SAGA components, depending whether glucose or ethanol is the carbon source. *GDH1*  
10 expression requires Ada2p and Ada3p on ethanol as carbon source, but only Ada3p on  
11 glucose. In both cases, *GDH1* expression is *GCN5* independent (32). In contrast, *GDH3*  
12 expression and chromatin remodelling activities in its cognate promoter, which is only  
13 seen under carbon de-repressing conditions (ethanol), are dependent on the SAGA  
14 components *GCN5*, *ADA2*, *ADA3* and *SPT3*.

15 There is little information on SAGA function in filamentous fungi. Recent work in  
16 *Neurospora crassa* established that blue light-induced transcription of the early light-  
17 inducible genes *al-3* and *vvd* depends on increased acetylation of histone H3 at lysine  
18 K14 in the promoters of these genes (18). The authors showed that acetylation and  
19 transient gene activation requires the *N. crassa* *GCN5* homologue *ngf-1* and it was  
20 suggested that the specific activator, White Collar-1, is required for the recruitment of  
21 NGF-1 to *al-3* and *vvd* promoters.

22 We have recently described the relationship between the transcriptional activation of a  
23 number of *A. nidulans* promoters driving genes of primary metabolism and their  
24 chromatin architecture (3, 12, 25, 27-29). The *prn* cluster of *A. nidulans* comprises all

1 the genes involved in proline utilisation as sole nitrogen and/or carbon source. We have  
2 studied in detail the bidirectional *prnD-prnB* promoter, driving respectively the genes  
3 encoding proline oxidase and the specific proline transporter (10, 11, 13-15, 34). The  
4 proline metabolic genes are induced by proline and repressed when preferred carbon  
5 (glucose) or nitrogen (ammonium, glutamine) sources are available in the medium. A  
6 graphical representation of regulatory proteins and their *cis*-acting recognition motifs  
7 involved in the regulation of the bidirectional promoter is shown in Figure 1A. In this  
8 promoter eight nucleosomes lose their positioning upon induction while simultaneous  
9 carbon and nitrogen metabolite repression results in partial nucleosome repositioning.  
10 Chromatin re-structuring is strictly dependent on the PrnA pathway-specific activator  
11 but not on the wide-domain nitrogen regulator AreA which was proven to be essential  
12 for chromatin opening in the *A. nidulans* nitrate regulon (3, 27). AreA in the proline  
13 cluster is only required in the presence of glucose, which is consistent with its proposed  
14 function which is to counteract the repressive effect of the global carbon repressor  
15 CreA. A functional CreA protein and two CreA binding sites in the *prnD-prnB*  
16 bidirectional promoter were shown to be essential for nucleosome re-positioning under  
17 repression conditions (12). A first indication that histone acetylation is involved in the  
18 CreA-mediated repression of the *prnD-prnB* bidirectional promoter process came from  
19 the observation that Trichostatin A, an inhibitor of histone deacetylases, mimics the  
20 effect of a CreA loss-of-function mutation or of *cis*-acting CreA binding site mutations  
21 (termed *prn<sup>d</sup>*). Nucleosome positioning after repression and transcriptional repression is  
22 completely lost when CreA is non functional and also partially lost when TSA is present  
23 in the culture medium (12).

1 To investigate in more detail how acetylation affects nucleosome positioning in this  
2 system, we deleted two genes coding for members of the putative SAGA/ADA  
3 acetylation complexes in *A. nidulans*. In this article we describe the effect of *gcnE* and  
4 *adaB* deletions, homologues of the *S. cerevisiae* *GCN5* and of *ADA2* genes, a putative  
5 histone acetylase and adaptor protein, respectively. We studied the effects of these  
6 deletions on the transcriptional competence, the chromatin structure and the acetylation  
7 status of histone H3 lysine K9 and K14 in the *prnD-prnB* bidirectional promoter.  
8 Unexpectedly, low levels of histone H3 acetylation in the deletion strains were found  
9 not to affect transcriptional activation, but are paradoxically associated with partial  
10 derepression of *prnB* and *prnD*.

11

## 12 **Materials and Methods**

### 13 **Strains and growth conditions.**

14 *A. nidulans* strains used throughout the study are listed in Table 1. A total of  $10^6$  spores  
15 of each strain per ml were inoculated into liquid minimal medium (30) with the  
16 appropriate supplements plus 0.1% fructose as the carbon source and 5 mM urea as the  
17 nitrogen source. Mycelia were grown for 12 h at 37°C on a rotary shaker with  
18 continuous shaking at 180rpm and then harvested by filtration. For condition-specific  
19 growth, aliquots of these pre-cultures were then further incubated for two hours at 37°C  
20 under following conditions: non induced, NI: 0,1% fructose, 5mM urea; induced, I: 20  
21 mM proline; induced repressed, IR: 20 mM proline plus 1% glucose and 20 mM  
22 ammonium-L[+]-tartrate. Finally, mycelia were harvested by filtration for RNA  
23 isolation, MNase analysis and chromatin immunoprecipitation. For microscopic  
24 observations, cultures were grown on solid complete media (30) until sporulation

1 occurred (roughly 7-10 days for the *adaB* and *gcnE* deletion strains and around 3 days  
2 for the wild type control strains) and surface samples were prepared for microscopy.

3

4

#### 5 **Cloning of *gcnE* and *adaB* and construction of disruption vector.**

6 Using the *S.cerevisiae* Gcn5p protein in a BLAST search of the *A.nidulans* genome  
7 database ([www.broad.mit.edu](http://www.broad.mit.edu)) we identified gene AN3621.3 as GcnE, a putative  
8 orthologue of yeast Gcn5p with an overall identity of 66% between the *A. nidulans* and  
9 the *S. cerevisiae* proteins (for predicted domain organisation and BLAST results, see  
10 [http://www.broad.mit.edu/annotation/genome/aspergillus\\_nidulans/](http://www.broad.mit.edu/annotation/genome/aspergillus_nidulans/)).

11 A deletion cassette was constructed by DJ-PCR. Deletion of the *gcnE* open reading  
12 frame (ORF) was achieved by replacing it with the *A. fumigatus pyrG* gene. In this  
13 cassette, the *pyrG* gene is flanked by upstream and downstream *gcnE* ORF sequences.

14 The 3' flanking fragment of *gcnE* was amplified with primers G1F (5'  
15 TAGCAGACCCTGATGCATCAAAC 3') and G2632R (5'  
16 TAAAGAGCTTTCAGTAAGTAATGATACTCG 3'), and the 5' flanking fragment of  
17 *gcnE* was amplified with G3330F (5'

18 GAAACATCCCCGAGTGGTCGGTAAGTACAA 3') and G5950R (5'  
19 CTCTCAAATACTAAGCCAAGTGGTACCAAG 3'), from *A. nidulans* wild type

20 (*pabaA1*) genomic DNA. *A. fumigatus pyrG* was amplified from *A. fumigatus* wild type  
21 genomic DNA with primers GCNPYRF (5'

22 GCGTGCTACCTCTGCGAGTATCATTACTTACTGAAAGCTCTTTAGGACTGAA

23 TTCGCCTCAAACAATGC 3') and GCNPYRR (5'

24 ATTGTACTTACCGACCACTGGGGATGTTTCGAATCTGCTGCCACATGAAGGA

1 ATTCTCAGTCCTGCTCC 3'). Nested primers GCNESTF (5'  
2 GAAAATTCAACTGTTCGATTCC 3') and GCNESTR (5'  
3 TGGGTGAAAAGTGGATTGTATG 3') were used to amplify the complete assembled  
4 molecule containing 5' *gcnE*-*pyrG*-3' *gcnE*. This *gcnE* deletion cassette was used to  
5 transform a *pyrG89 argB2 pantoB100 riboB2 yA2* strain. Transformants were selected  
6 on minimal media with appropriate supplements lacking uracil and uridine.

7 To identify the strains deleted for *gcnE*, a Southern blot with genomic DNA digested  
8 with *KpnI* was hybridized with a <sup>32</sup>P labeled probe derived from a 2,6 kb PCR product  
9 amplified with primers G3330F and G5950R. A single 3 kb band confirmed *gcnE*  
10 deletion and single integration of the deletion cassette. The same membrane was  
11 stripped and hybridized with a <sup>32</sup>P labeled probe derived from a 800 pb PCR product  
12 amplified by GCNORFF (5' TCGTTGGAGGTATAACCTACCG 3') and GCNORFR  
13 (5' GTAGGGTGTGTTCTCGTTATTATACC 3') corresponding to the *gcnE* ORF. No  
14 signal was detected for the *gcnE* deleted strain (not shown).

15 Using the *S.cerevisiae* Ada2p protein in a BLAST search of the *A.nidulans* genome  
16 database ([www.broad.mit.edu](http://www.broad.mit.edu)) we identified gene AN5974.2 (changed to gene number  
17 AN10763.3 in the latest annotation) as a putative orthologue of yeast Ada2p with a  
18 similarity score of  $3 \times 10^{-79}$  (for predicted domain organisation and BLAST results, see  
19 [http://www.broad.mit.edu/annotation/genome/aspergillus\\_nidulans/](http://www.broad.mit.edu/annotation/genome/aspergillus_nidulans/)). Transcription  
20 start and termination points were determined empirically by 5' and 3' RACE,  
21 respectively, and were found to correspond to the sites shown in the *A. nidulans*  
22 database. Plasmid pMS12 (Fungal Genetics Stock Center, <http://www.fgsc.net>)  
23 containing the *argB* gene was used to insert PCR fragments derived from the *adaB* 5'  
24 region upstream of *argB* and from the *adaB* 3' region downstream of *argB*. The *adaB*



1 5' and 3' sequences were obtained by PCR amplification using the sequence information  
2 from the *A.nidulans* sequence database ([www.broad.mit.edu](http://www.broad.mit.edu)). Roughly 700 bp of the  
3 *adaB* 5' region was amplified with primers 5' forward:  
4 5'GATGCAAGACCGCGGCCGCCACAGCAGTCAGC 3' (NotI site introduced by  
5 mutating three bases is underlined) and 5' reverse:  
6 5'CATCGATCCCGAGAGCCTTTCATCCTCCGCG 3'). The fragment was cloned as  
7 NotI-ClaI fragment into the vector. Around 1,4 kb of the *adaB* 3' region was amplified  
8 with primers 3' forward: 5'CGAATCATCACGTCGACTTGAATTTAGAGGCC 3'  
9 (SalI site introduced by mutation of the HindIII restriction site, positioned 50 bp  
10 downstream of the empirically determined poly adenylation site, is underlined) and  
11 3' reverse: 5'GCGAGTTGACTGAGCTCGAGAAGGATCACCCCTC 3'. The fragment  
12 was cloned as Sal I-Xho I fragment into the vector. The resulting 7,2 kb vector was  
13 called p5'-3'adaB::argB. Strain MH 9233 (26) (gift from Michael Hynes, The  
14 University of Melbourne) was used as recipient strain for the knock out construct. To  
15 verify the insertion of the *argB* gene at the *adaB* locus, Southern hybridisation was  
16 carried out and the putative deletion strains (showing absence of the 4,5 kb *adaB* band  
17 but instead, a single 3,5 kb band hybridizing with the 3'*adaB* Hind III-Xho I probe)  
18 were verified for the lack of the *adaB* ORF and additional *argB* insertions. The strain  
19 was crossed to a wild type *pabaA1* strain to obtain an *adaB* $\Delta$  strain with only one  
20 additional marker for further studies. *adaB* $\Delta$  *biA1* was recovered from the progeny and  
21 for further studies, a wild type *biA1* strain was used as reference strain.

22 **RNA preparation and Northern blots.** Total RNA was isolated with the RNA Plus  
23 Extraction Solution (Biogen) following the manufacturer's instructions. RNA

1 electrophoresis and Northern blot hybridizations were carried out as described  
2 previously (13). *prnB*, *prnD*, and *acnA* probes were prepared as described by (14).

3

4 **Nucleosome positioning.** Micrococcal nuclease I digestions were performed by the  
5 method adapted by Gonzalez *et al* (16). Micrococcal nuclease was used at  
6 concentrations ranging between 0.5 and 2.5 U/g of mycelium. DNA was digested with  
7 an appropriate restriction enzyme. Probes SC1 and SC2 for the *prnB-D* intergenic  
8 region were prepared as described previously (12)

9 The position of each MNase cut was calculated by running in each gel a scale of  
10 molecular size markers (100 Base-Pair Ladder, Amersham Pharmacia Biotech.,  
11 Piscataway, NJ). The values reported in the scale adjacent to each gel represent the  
12 position of the MNase cut from the ATG of the relevant gene. For each mutant and  
13 growth condition the experiments were performed in triplicate.

14

#### 15 **ChIP assays**

16 Chromatin immunoprecipitation assays (ChIP) were carried out following our published  
17 protocol (4). Antibodies for ChIP analysis of acetylated histone H3 were purchased  
18 from Upstate Biotechnology, Charlottesville, VA, recognizing acetylated K9 and K14  
19 of histone H3 (#06-599). Rabbit polyclonal antibody recognizing the C-terminus of  
20 histone H3 (#ab1791) was purchased from abcam (Cambridge, UK). Latter antibody  
21 was used to normalize the acetylation status of H3. 2 µl of antibody solution were used  
22 to incubate 200 µg of protein.

23 Quantitative real-time PCR used a BIO-RAD (Hercules, CA) MyiQ Cyclyer with the  
24 Platinum® SYBR® Green qPCR SuperMix-UDG from Invitrogen (Karlsruhe,

1 Germany) for amplification. Chromosomal DNA from a wild-type strain was used as  
2 external standard for setting up the calibration curve. Primers which amplified  
3 Fragment **a** (see Figure 4B) had the following sequence: prnB+2for:  
4 5'TGAGGATCCCATTAGTCAAGG3' and prnB+2rev:  
5 5'GGATCAGGTTCCCTAAGATCAG3'. PCR was calibrated by a dilution series of  
6 total DNA extracted from a wild-type strain (3.3  $\mu\text{g } \mu\text{l}^{-1}$ ). For each sample the absolute  
7 amount of the specific DNA fragment in the immunoprecipitated sample was divided by  
8 the amount of this fragment in the sample before precipitation. Each PCR reaction was  
9 replicated (technical repetition). The resulting ratio of the precipitation with antibody  
10 H3-acetyl was divided by the resulting ratio of the antibody H3 C-term precipitation and  
11 the resulting ratio of the WT strain treated for non inducing conditions was set to one.  
12 Two biological repetitions were performed for each condition and standard deviations  
13 were calculated.

## 15 **Results**

### 16 **Phenotypes of the *gcnE* and *adaB* deletions**

17 Both the *gcnE* and *adaB* deletion strains show a strongly reduced growth rate and  
18 conidiation on solid medium. Both deletions resulted in similar morphological  
19 alterations of the conidiophore. Supplementary Figure S1 shows a selection of pictures  
20 obtained from microscopic observations of the *adaB* deletion strain. The strain produces  
21 stalks and vesicles, but metulae are missing. Short stalks with vesicle heads and a few  
22 phialides repeatedly emerge directly from vegetative mycelia. None of the known  
23 conidiation mutants of *A. nidulans* displays this phenotype.

1 We tested for specific effects of the *gcnEΔ* and *adaBΔ* strains on the utilization of  
2 various carbon and nitrogen sources. Compared to the isogenic wild type strains, both  
3 deletion strains displayed a reduced growth on all the nutrients tested, including  
4 complete media and proline as sole carbon or nitrogen source (data not shown).

5

6 ***prnD-prnB* expression is partially de-repressed in *gcnEΔ* and *adaBΔ*.**

7 García and co-workers (2004) have shown that expression of proline catabolic genes  
8 and chromatin rearrangements in the *prnD-prnB* intergenic region are affected by  
9 trichostatin-A mediated inhibition of deacetylases. Thus we investigated expression of  
10 *prnB* and *prnD* in a *gcnEΔ* and *adaBΔ* background. Figure 1B shows Northern blots  
11 from isogenic *gcnE<sup>+</sup>* and *gcnEΔ* (right panel) as well as isogenic *adaB<sup>+</sup>* and *adaBΔ*  
12 strains (left panel) grown under non-inducing (NI, fructose, urea), inducing (I, proline)  
13 and inducing-repressing (IR, proline, glucose, ammonium) conditions. The results of  
14 deleting *gcnE* or *adaB* on the expression of *prnB* and *prnD* are surprising. There is no  
15 effect on the non-induced (undetectable) and induced levels. However, a partial de-  
16 repression of *prnB* and *prnD* is seen in both deletions strains. Previous work (10) has  
17 shown that only the repression of *prnB* is direct, the repression of *prnD* being the result  
18 of inducer exclusion. The latter easily accounts for the fact that the mild de-repression  
19 of *prnB* results in an elevated de-repression of *prnD*, particularly pronounced in the  
20 *gcnEΔ* strain. Interestingly, neither deletion significantly alters the induced levels of  
21 *prnB* or *prnD* transcripts, suggesting that the induction process could be independent of  
22 the histone acetylation status.

23

1 **Chromatin organisation in the *prnB-prnD* promoter is affected in SAGA/ADA**  
2 **component deletion strains.**

3 Figure 2A shows the results of a micrococcal nuclease (MNase) digestion assay for  
4 *adaB*<sup>+</sup> and *adaB*Δ under the three standard conditions investigated. The results of  
5 MNase digestions in the *gcnE*Δ strain (not shown) are identical to those obtained for the  
6 *adaB* deletion. In both deletion strains, the only difference to the corresponding isogenic  
7 wild type control in the nuclease digestion pattern is observed under induced-repressed  
8 conditions (IR). In an *adaB*<sup>+</sup> *gcnE*<sup>+</sup> strain all nucleosomes are re-positioned under IR  
9 conditions (12), whereas in the *adaB*Δ and *gcnE*Δ strains, this re-positioning is  
10 defective. In the deletion strains the IR nucleosome pattern is identical to that seen  
11 under induced conditions which is also identical to the pattern obtained under induced  
12 conditions in the wild type strain. Hence, the derepressed expression of *prnB* and *prnD*  
13 observed in the *adaB* and *gcnE* deleted strains is associated with and could depend on  
14 defective nucleosome re-positioning. The SC1 probe shown here allows the detection of  
15 the region including from nucleosome -4 to nucleosome +2. The probe detecting the  
16 region including nucleosomes +3 and +4 (probe SC2) did not reveal any differences  
17 between the deletion strains and the corresponding wild type strains (data not shown). A  
18 graphical representation of nucleosome positioning in the *prnB-prnD* promoter region is  
19 shown in Figure 2B. Nucleosome positioning in the *adaB*Δ strain under IR conditions  
20 and a de-repressed phenotype for *prnB* and *prnD* expression closely resembles results  
21 obtained when the de-acetylase inhibitor trichostatin A is used in the wild type or when  
22 MNase patterns are analysed in a *creA*<sup>d1</sup> strain (10). We thus used ChIP to determine  
23 the acetylation status of histone H3 in the region of nucleosome +2, the only

1 nucleosome that remains positioned under these conditions in the *adaBA* and *gcnEA*  
2 strains.

3

4 **The acetylation status of histone H3 in the region of nucleosome +2 depends on**  
5 **physiological conditions and on the function of GcnE and AdaB.**

6 We chose to look at the region of nucleosome +2, as MNase digests indicate that a  
7 nucleosome is present in this region under all conditions, whether completely positioned  
8 (non induced conditions) or partially positioned (induced and induced repressed  
9 conditions) (12). The acetylation status of lysines 9 (H3K9) and 14 (H3K14) in histone  
10 H3 is a critical mark for gene activation. To investigate the acetylation status of H3K9  
11 and H3K14 in nucleosome +2 we employed chromatin immunoprecipitation (ChIP)  
12 assays using an antibody recognising both H3K9ac and H3K14ac followed by  
13 amplification of a fragment encompassing the region covered by nucleosome +2  
14 (fragment **a** in Figure 4B). The ratio between acetylated and total H3 (precipitated with  
15 an antibody directed against a H3 C-terminal epitope) is shown in Figure 3. Both  
16 acetylation patterns in *adaBA* and *gcnEA* strains are very similar, but, in contrast to the  
17 MNase digests, they are not identical and are therefore presented here for both strains.

18 In the *adaB*<sup>+</sup> (wild type) strain, isogenic to the *adaBA* strain, acetylation of K9 and K14  
19 in H3 increases with induction indicating a correlation between H3 acetylation,  
20 chromatin rearrangements (loss of positioning) and transcriptional activation. Upon  
21 repression, the acetylation status of H3K9 and H3K14 does not change significantly,  
22 which indicates that de-acetylation of these residues in H3 is not required to trigger gene  
23 repression and re-positioning of nucleosome +2. In contrast to *adaB*<sup>+</sup>, in the *adaBA*  
24 strain, acetylation of H3K9 and K14 is strongly reduced under all conditions and despite

1 considerable induction of *prnB* and *prnD* transcription, an increase in acetylation of H3  
2 is not seen in these strains. A qualitatively very similar pattern is observed when the  
3 *gcnE*<sup>+</sup> (wild type) control strain is compared to the isogenic *gcnEΔ* strain under the  
4 different growth conditions. As in the *adaBΔ* strain, acetylation of H3K9 and K14 is  
5 strongly reduced in the *gcnE* deletion strain under all conditions.

6

## 7 **Discussion**

8 Often, enhanced gene expression correlates with loss of positioning and increased  
9 acetylation of nucleosomes (33). In the *prnD-prnB* bidirectional promoter, we see  
10 increased acetylation of lysines 9 and 14 in histone H3 under inducing conditions in the  
11 region corresponding to partially positioned nucleosome +2. However, this increased  
12 acetylation is not required for efficient transcription or chromatin remodelling. In the  
13 strains lacking essential components of the putative SAGA/ADA complexes, i.e. GcnE  
14 or AdaB, we detect only low levels of H3 lysine acetylation, but induction and  
15 remodelling of chromatin are not affected. This is not without precedence as in *S.*  
16 *cerevisiae*, SAGA/ADA-dependent and independent activators are known. Activation of  
17 transcription requires SAGA or ADA complexes in the GAL1 promoter where GAL4  
18 recruits the SAGA complex, and this in turn is necessary for the recruitment of the  
19 mediator complex (20). Our results indicate that, while a putative SAGA or ADA  
20 complex might be involved in determining the acetylation levels in this region, PrnA-  
21 mediated activation and chromatin remodelling of the *prnD-prnB* promoter does not  
22 depend on AdaB and GcnE, putative members of *A. nidulans* SAGA/ADA complexes.

23 It should be noted that in our ChIP experiments we have only tested histone H3  
24 acetylation at lysines 9 and 14, residues known to be acetylated by SAGA and ADA

1 complexes (9). Other covalent modifications not detected by the antibody used here,  
2 such as acetylations of H2A, H2B and H4 at diverse lysines in their N-termini could  
3 also lead to hyperacetylation of nucleosomes in the *prnD-prnB* promoter thus providing  
4 the signal for effective chromatin remodelling. In filamentous fungi, it is completely  
5 unknown which mechanisms direct the cross-talk of histone modification. In a study  
6 focussing on the different roles of H3 and H4 acetylation in the yeast *Saccharomyces*  
7 *cerevisiae*, it was shown that H4 hyperacetylation in promoters of Adr1p-dependent  
8 genes does not compensate for low H3 acetylation in a *gcn5* mutant strain (1).  
9 The observed partial de-repression of *prnB* and *prnD* expression in the *adaB* and *gcnE*  
10 deleted strains was unexpected. However, in *Neurospora crassa*, mutation of the GCN5  
11 homologue *ngf-1* results in loss of repression of the blue light-responsive *al-3* gene in  
12 the dark (18). A *ngf-1<sup>RIP1</sup>* mutant was found to express *al-3* constitutively suggesting  
13 that the histone acetyltransferase is not only involved in blue-light induction but also in  
14 repression of *al-3* in the absence of light. Another example of HAT activities being  
15 involved in both, gene activation and repression is provided by the yeast GAL system.  
16 Glucose repression requires ADA2p and ADA3p/NGG1p, an ADA-complex protein  
17 functionally associated with GCNp5 and ADA2p (6) and it has been suggested that  
18 ADA2p/ADA3p directly inhibits the activation domain of GAL4p. Similarly, in the  
19 yeast arginine metabolic system GCN5 function and increased histone acetylation are  
20 associated with transcriptional repression. In this case, Gcn5p-mediated increase of H3  
21 acetylation correlates with binding of the arginine repressor complex ARGR/MCM1  
22 (31).  
23 However, our nucleosome positioning and ChIP studies lead to a different picture.  
24 Although histone H3 acetylation increases during transcriptional activation, it is neither



1 required for transcriptional activation (compare induced conditions in *gcnE<sup>+</sup>adaB<sup>+</sup>* with  
2 *gcnEΔ* or *adaBΔ* in Figures 1 and 3) nor loss of nucleosome positioning (compare  
3 Figure 2). Additionally, transcriptional repression and re-positioning of nucleosomes are  
4 not accompanied by de-acetylation of histone H3 at K9 or K14 (see Figure 3, IR  
5 conditions). These results suggest that differences in histone acetylation are associated  
6 with, but not necessary for transcriptional activation and nucleosome positioning  
7 processes in the *prnD-prnB* intergenic region. It was therefore surprising to see that  
8 both, hyper-acetylation by TSA inhibition of deacetylases and hypoacetylation by  
9 deletion of *adaB* or *gcnE* leads to partial de-repression of the proline catabolic genes  
10 and simultaneous lack of nucleosome re-positioning.

11 Transcriptional repression and nucleosome re-positioning upon simultaneous glucose  
12 and ammonium repression requires a functional CreA protein. This has been shown by  
13 mutations in both the CreA protein itself (*creA<sup>d1</sup>*) and in its cognate binding sites  
14 (*prn<sup>d20</sup>* or *prn<sup>d22</sup>*) in the *prnD-prnB* promoter (12). The simplest hypothesis integrating  
15 our experimental results would therefore propose that both, hyperacetylation by TSA  
16 treatment and hypoacetylation by *adaB* or *gcnE* deletion impair CreA function. Several  
17 observations support this view. Firstly, nucleosomal patterns obtained in the *prnD-prnB*  
18 region by TSA treatment and in *adaBΔ* and *gcnEΔ* under IR conditions are identical to  
19 those found in *creA<sup>d1</sup>*, *prn<sup>d20</sup>* or *prn<sup>d22</sup>* mutant strains (13). Secondly, de-repression  
20 under TSA treatment or in *adaBΔ* and *gcnEΔ* is only partial whereas de-repression is  
21 complete in *creA* loss-of-function strains. This also implies that while nucleosome re-  
22 positioning may be necessary for full repression, CreA can still partially repress on  
23 completely open chromatin, probably by directly interfering with the activating function  
24 of the pathway-specific activator PrnA.

1 Thirdly, the function of CreA, but not the acetylation status of histones, defines whether  
2 nucleosomes are positioned or not. In summary, nucleosomal rearrangements,  
3 transcriptional activity and histone acetylation are clearly distinct processes in the *prnD*-  
4 *prnB* region and therefore a direct influence of the acetylation status on nucleosomal  
5 positioning and transcription is highly unlikely. Further studies will reveal at which  
6 level, transcriptional and/or post-transcriptional, CreA function is impaired by TSA  
7 treatment or lack of histone acetyl transferase activities.

8

### 9 **Acknowledgements**

10 We are grateful for receiving strain MH9233 from Michael Hynes (The University of  
11 Melbourne). The work was supported by grant number P17018 of the Austrian Science  
12 Fund FWF to JS. Work at Orsay was supported by the CNRS, the Université Paris-Sud,  
13 and the Institut Universitaire de France. Y.R.-D. was supported by a predoctoral  
14 studentship of the Ministère de l'Éducation Supérieure et de la Recherche and the  
15 CONACYT (México). R.F.-M. was supported by a Marie Curie Fellowship (MCFI-  
16 2001-01084). I.G. was supported by postdoctoral fellowships from, successively, the  
17 Spanish Ministry of Education and the EU Marie Curie Programme.

1

2 **References**

- 3 1. **Agricola, E., L. Verdone, E. Di Mauro, and M. Caserta.** 2006. H4 acetylation  
4 does not replace H3 acetylation in chromatin remodelling and transcription  
5 activation of Adr1-dependent genes. *Mol Microbiol* **62**:1433-46.
- 6 2. **Balasubramanian, R., M. G. Pray-Grant, W. Selleck, P. A. Grant, and S.**  
7 **Tan.** 2002. Role of the Ada2 and Ada3 transcriptional coactivators in histone  
8 acetylation. *J Biol Chem* **277**:7989-95.
- 9 3. **Berger, H., R. Pachlinger, I. Morozov, S. Goller, F. Narendja, M. Caddick,**  
10 **and J. Strauss.** 2006. The GATA factor AreA regulates localization and in vivo  
11 binding site occupancy of the nitrate activator NirA. *Mol. Microbiol.* **59**:433-  
12 446.
- 13 4. **Bernreiter, A., A. Ramon, J. Fernandez-Martinez, H. Berger, L. Araujo-**  
14 **Bazan, E. A. Espeso, R. Pachlinger, A. Gallmetzer, I. Anderl, C.**  
15 **Scazzocchio, and J. Strauss.** 2007. Nuclear export of the transcription factor  
16 NirA is a regulatory checkpoint for nitrate induction in *Aspergillus nidulans*.  
17 *Mol Cell Biol* **27**:791-802.
- 18 5. **Bhaumik, S. R., and M. R. Green.** 2001. SAGA is an essential in vivo target of  
19 the yeast acidic activator Gal4p. *Genes Dev* **15**:1935-45.
- 20 6. **Brandl, C. J., J. A. Martens, A. Margaliot, D. Stenning, A. M. Furlanetto,**  
21 **A. Saleh, K. S. Hamilton, and J. Genereaux.** 1996. Structure/functional  
22 properties of the yeast dual regulator protein NGG1 that are required for glucose  
23 repression. *J Biol Chem* **271**:9298-306.
- 24 7. **Brownell, J. E., J. Zhou, T. Ranalli, R. Kobayashi, D. G. Edmondson, S. Y.**  
25 **Roth, and C. D. Allis.** 1996. *Tetrahymena* histone acetyltransferase A: a

- 1 homolog to yeast Gcn5p linking histone acetylation to gene activation. Cell  
2 **84**:843-51.
- 3 8. **Candau, R., J. X. Zhou, C. D. Allis, and S. L. Berger.** 1997. Histone  
4 acetyltransferase activity and interaction with ADA2 are critical for GCN5  
5 function in vivo. *Embo J* **16**:555-65.
- 6 9. **Carrozza, M. J., R. T. Utley, J. L. Workman, and J. Cote.** 2003. The diverse  
7 functions of histone acetyltransferase complexes. *Trends Genet* **19**:321-9.
- 8 10. **Cubero, B., D. Gomez, and C. Scazzocchio.** 2000. Metabolite repression and  
9 inducer exclusion in the proline utilization gene cluster of *Aspergillus nidulans*.  
10 *J Bacteriol* **182**:233-5.
- 11 11. **Cubero, B., and C. Scazzocchio.** 1994. Two different, adjacent and divergent  
12 zinc finger binding sites are necessary for CREA-mediated carbon catabolite  
13 repression in the proline gene cluster of *Aspergillus nidulans*. *Embo J* **13**:407-  
14 15.
- 15 12. **Garcia, I., R. Gonzalez, D. Gomez, and C. Scazzocchio.** 2004. Chromatin  
16 rearrangements in the *prnD-prnB* bidirectional promoter: dependence on  
17 transcription factors. *Eukaryot Cell* **3**:144-56.
- 18 13. **Gomez, D., B. Cubero, G. Cecchetto, and C. Scazzocchio.** 2002. PrnA, a  
19 Zn<sub>2</sub>Cys<sub>6</sub> activator with a unique DNA recognition mode, requires inducer for in  
20 vivo binding. *Mol Microbiol* **44**:585-97.
- 21 14. **Gomez, D., I. Garcia, C. Scazzocchio, and B. Cubero.** 2003. Multiple GATA  
22 sites: protein binding and physiological relevance for the regulation of the  
23 proline transporter gene of *Aspergillus nidulans*. *Mol Microbiol* **50**:277-89.

- 1 15. **Gonzalez, R., V. Gavrias, D. Gomez, C. Scazzocchio, and B. Cubero.** 1997.  
2 The integration of nitrogen and carbon catabolite repression in *Aspergillus*  
3 *nidulans* requires the GATA factor AreA and an additional positive-acting  
4 element, ADA. *Embo J* **16**:2937-44.
- 5 16. **Gonzalez, R., and C. Scazzocchio.** 1997. A rapid method for chromatin  
6 structure analysis in the filamentous fungus *Aspergillus nidulans*. *Nucleic Acids*  
7 *Res* **25**:3955-6.
- 8 17. **Grant, P. A., L. Duggan, J. Cote, S. M. Roberts, J. E. Brownell, R. Candau,**  
9 **R. Ohba, T. Owen-Hughes, C. D. Allis, F. Winston, S. L. Berger, and J. L.**  
10 **Workman.** 1997. Yeast Gcn5 functions in two multisubunit complexes to  
11 acetylate nucleosomal histones: characterization of an Ada complex and the  
12 SAGA (Spt/Ada) complex. *Genes Dev* **11**:1640-50.
- 13 18. **Grimaldi, B., P. Coiro, P. Filetici, E. Berge, J. R. Dobosy, M. Freitag, E. U.**  
14 **Selker, and P. Ballario.** 2006. The *Neurospora crassa* White Collar-1  
15 dependent blue light response requires acetylation of histone H3 lysine 14 by  
16 NGF-1. *Mol Biol Cell* **17**:4576-83.
- 17 19. **Koehler, R. N., N. Rachfall, and R. J. Rolfes.** 2007. Activation of the ADE  
18 genes requires the chromatin remodeling complexes SAGA and SWI/SNF.  
19 *Eukaryot Cell* **6**:1474-85.
- 20 20. **Koh, S. S., A. Z. Ansari, M. Ptashne, and R. A. Young.** 1998. An activator  
21 target in the RNA polymerase II holoenzyme. *Mol Cell* **1**:895-904.
- 22 21. **Kuo, M. H., J. E. Brownell, R. E. Sobel, T. A. Ranalli, R. G. Cook, D. G.**  
23 **Edmondson, S. Y. Roth, and C. D. Allis.** 1996. Transcription-linked

- 1 acetylation by Gcn5p of histones H3 and H4 at specific lysines. *Nature* **383**:269-  
2 72.
- 3 22. **Larschan, E., and F. Winston.** 2001. The *S. cerevisiae* SAGA complex  
4 functions in vivo as a coactivator for transcriptional activation by Gal4. *Genes*  
5 *Dev* **15**:1946-56.
- 6 23. **Lee, T. I., H. C. Causton, F. C. Holstege, W. C. Shen, N. Hannett, E. G.**  
7 **Jennings, F. Winston, M. R. Green, and R. A. Young.** 2000. Redundant roles  
8 for the TFIID and SAGA complexes in global transcription. *Nature* **405**:701-4.
- 9 24. **Lee, T. I., and R. A. Young.** 2000. Transcription of eukaryotic protein-coding  
10 genes. *Annu Rev Genet* **34**:77-137.
- 11 25. **Mathieu, M., I. Nikolaev, C. Scazzocchio, and B. Felenbok.** 2005. Patterns of  
12 nucleosomal organization in the *alc* regulon of *Aspergillus nidulans*: roles of the  
13 AlcR transcriptional activator and the CreA global repressor. *Mol Microbiol*  
14 **56**:535-48.
- 15 26. **Monahan, B. J., J. A. Fraser, M. J. Hynes, and M. A. Davis.** 2002. Isolation  
16 and characterization of two ammonium permease genes, *meaA* and *mepA*, from  
17 *Aspergillus nidulans*. *Eukaryot Cell* **1**:85-94.
- 18 27. **Muro-Pastor, M. I., R. Gonzalez, J. Strauss, F. Narendja, and C.**  
19 **Scazzocchio.** 1999. The GATA factor AreA is essential for chromatin  
20 remodelling in a eukaryotic bidirectional promoter [published erratum appears in  
21 *EMBO J* 1999 May 4;18(9):2670]. *Embo J* **18**:1584-97.
- 22 28. **Muro-Pastor, M. I., J. Strauss, A. Ramon, and C. Scazzocchio.** 2004. A  
23 paradoxical mutant GATA factor. *Eukaryot Cell* **3**:393-405.

- 1 29. **Narendja, F., S. P. Goller, M. Wolschek, and J. Strauss.** 2002. Nitrate and the  
2 GATA factor AreA are necessary for in vivo binding of NirA, the pathway-  
3 specific transcriptional activator of *Aspergillus nidulans*. *Mol Microbiol* **44**:573-  
4 83.
- 5 30. **Pontecorvo, G., J. A. Roper, L. M. Hemmons, K. D. MacDonald, and A. W.**  
6 **J. Bufton.** 1953. The genetics of *Aspergillus nidulans*. *Adv. Genet.* **5**:141-238.
- 7 31. **Ricci, A. R., J. Genereaux, and C. J. Brandl.** 2002. Components of the SAGA  
8 histone acetyltransferase complex are required for repressed transcription of  
9 ARG1 in rich medium. *Mol Cell Biol* **22**:4033-42.
- 10 32. **Riego, L., A. Avendano, A. DeLuna, E. Rodriguez, and A. Gonzalez.** 2002.  
11 GDH1 expression is regulated by GLN3, GCN4, and HAP4 under respiratory  
12 growth. *Biochem Biophys Res Commun* **293**:79-85.
- 13 33. **Roth, S. Y., J. M. Denu, and C. D. Allis.** 2001. Histone acetyltransferases.  
14 *Annu Rev Biochem* **70**:81-120.
- 15 34. **Sophianopoulou, V., and C. Scazzocchio.** 1989. The proline transport protein  
16 of *Aspergillus nidulans* is very similar to amino acid transporters of  
17 *Saccharomyces cerevisiae*. *Mol Microbiol* **3**:705-14.
- 18 35. **Wang, L., C. Mizzen, C. Ying, R. Candau, N. Barlev, J. Brownell, C. D.**  
19 **Allis, and S. L. Berger.** 1997. Histone acetyltransferase activity is conserved  
20 between yeast and human GCN5 and is required for complementation of growth  
21 and transcriptional activation. *Mol Cell Biol* **17**:519-27.
- 22  
23

1 Table 1: *Aspergillus nidulans* strains used throughout this work

Strain	Genotype	Reference
<i>gcnE</i> <sup>+</sup>	<i>pyrG89 argB2 pantoB100</i> <i>riboB2 yA2</i>	This work
<i>gcnEΔ</i>	<i>pyrG89 argB2 pantoB100</i> <i>riboB2 yA2 gcnEΔ::pyrG</i> <sup>+</sup>	This work
<i>adaB</i> <sup>+</sup>	<i>biA1</i>	FGSC <sup>1</sup> strain #A26
<i>adaBΔ</i>	<i>biA1 argB::trpCΔB</i> <i>adaBΔ::argB</i> <sup>+</sup>	This work
MH 9233	<i>wA3 biA1 argB::trpCΔB</i> <i>pyroA4 riboB2</i>	(26)

2 <sup>1</sup>Fungal Genetics Stock Center: <http://www.fgsc.net>



1

## 2 **Figure Legends**

3 **Figure 1.** Regulation of proline utilization in *A. nidulans*. **A.** Overview of proline  
4 cluster regulation. The *prnD-B* intergenic region is shown. *prnD* encodes the proline  
5 oxidase, and *prnB* encodes the specific proline transporter (12, 14). The pathway-  
6 specific transcription factor PrnA is essential for proline induction of both genes. In the  
7 absence of preferential carbon (glucose) and nitrogen (ammonium) sources and the  
8 presence of proline, PrnA and the GATA factor AreA bind to their cognate sites in the  
9 intergenic region resulting in the expression of *prnD* and *prnB*. Repression requires both  
10 glucose activation of the negative regulator CreA and ammonium inactivation of AreA.  
11 Full repression occurs only in the simultaneous presence of glucose and ammonium.  
12 Repression acts directly on *prnB* expression, *prnD* repression is indirect and results  
13 from inducer exclusion. Activation processes are represented by arrows and repression  
14 by terminals.

15 **B.** Effect of *adaB* and *gcnE* deletion on *prnB* and *prnD* transcription. Strains (see Table  
16 1) were pre-grown in liquid minimal medium under non-inducing conditions (5 mM  
17 urea, 0,1 % fructose) with the appropriate supplements, harvested, divided into aliquots  
18 und further incubated for two hours under conditions indicated. NI, 5 mM urea and  
19 0.1% fructose; I, induced by 20mM proline, IR, 20 mM proline and repression by 1%  
20 glucose and 20 mM ammonium-L(+)-tartrate (IR). Expression levels (bottom) were  
21 quantified using phosphoimaging and ImageQuant software analysis. Normalised  
22 signals were obtained by comparison of specific signals with actin gene (*acnA*)  
23 expression signals. The induced level in the *adaB*<sup>+</sup> and *gcnE*<sup>+</sup> strains is given in every

1 case the arbitrary value of 100, filled columns represent *prnB* and open columns *prnD*  
2 expression.

3


4 **Figure 2.** Nucleosome positioning in the *prnD-prnB* intergenic region in *adaB*<sup>+</sup> and  
5 *adaBΔ* strains. **A.** Indirect end-labelling micrococcal nuclease I (MNaseI) analysis of  
6 the *prnD-prnB* promoter region using probe SC1. This probe reveals nucleosomes +2 to  
7 -4. MNase analysis of nucleosomes +3 and +4 are not shown as the pattern obtained for  
8 the mutant strains are identical to the one extant in the wild type. Growth conditions  
9 (NI, I, IR) were identical to those described in Figure 3. Numbers besides the  
10 autoradiograms correspond to the positions of the main cuts relative to the *prnD* ATG.  
11 These were calculated from molecular size markers run in every gel. Asterisks indicate  
12 the position of the relevant changes observed. Triangles indicate increasing  
13 concentration of MNaseI. nDNA, naked DNA. To the left of the *adaBΔ* NI lanes a  
14 schematic representation of the nucleosome structure is shown. The wild type pattern  
15 under all conditions have been previously published (12), we only include in the same  
16 gel, for comparison, the pattern obtained under IR conditions, the only one where an  
17 *adaB* deletion pattern differs from that of the wild type

18 **B.** Schematic representation of nucleosome positioning (based on figure 2A and  
19 reference (12)). All eight nucleosomes of the intergenic region are drawn. Arrows  
20 indicate MNase I digests. Their thickness indicates the relative intensity of the bands in  
21 the autoradiogram shown in figure 2A and in other aoradiograms (not shown) covering  
22 nucleosomes +3 and +4. Dashed arrows indicate weakly cut sites. Under non-induced  
23 (NI) and induced (I) conditions, the nucleosome patterns are identical in strain *adaB*<sup>+</sup>  
24 and *adaBΔ*. White ovals represent fully positioned nucleosomes, while partially


1 positioned nucleosomes are shown by diagonally hatched ovals. The interpretation and  
2 significance of this partial positioning is discussed in the text and in references (12, 25).  
3 Symbols: white lozenges, CreA-binding sites 3.1 and 3.2, which are mutated in the  
4 *prn<sup>d</sup>20* and *prn<sup>d</sup>22* strains and result in derepression (see text); grey ovals, AreA-  
5 binding sites 13 and 14, shown to be the physiologically relevant AreA binding sites  
6 (15); white triangles, high-affinity PrnA binding sites 2 and 3 (14); black triangle,  
7 TATA box (15); The fragment amplified in the ChIP analysis (a) is shown as solid line  
8 below nucleosome +2. nDNA, pattern obtained with naked DNA, to the right of the  
9 scheme we indicate to which strain(s) a given nucleosome pattern correspond.


10  
11 **Figure 3.** Chromatin immunoprecipitation (ChIP) assay comparing acetylated H3 K9  
12 and K14 in nucleosome +2 of the *prnD-prnB* intergenic region between *adaB<sup>+</sup>* and  
13 *adaBΔ* (panel A) and between *gcnE<sup>+</sup>* and *gcnEΔ* (panel B) strains (see Table 1 for  
14 genotype details). The ratio between total H3 (C-terminal epitope) and H3-acetyl  
15 (recognizing H3 K9/K14) in the wild type strains (*adaB<sup>+</sup>* or *gcnE<sup>+</sup>*) grown under non-  
16 inducing conditions (NI) was set to 1. Error bars are indicating the standard deviation of  
17 four experiments (two biological and two technical repetitions per condition). Growth  
18 conditions: NI, non-inducing; I, inducing; IR, simultaneously inducing-repressing.

19




B






**B**



A





B

